

DISSERTATIONES SCHOLAE DOCTORALIS AD SANITATEM INVESTIGANDAM
UNIVERSITATIS HELSINKIENSIS

**SUSANNA JOUHI
NEE SALKOLA**

**SMALL CHOROIDAL MELANOMAS
— MANAGEMENT AND PROGNOSIS**



DEPARTMENT OF OPHTHALMOLOGY
HELSINKI UNIVERSITY HOSPITAL
AND
FACULTY OF MEDICINE
DOCTORAL PROGRAMME IN CLINICAL RESEARCH
UNIVERSITY OF HELSINKI

Department of Ophthalmology,
Helsinki University Hospital
and
Faculty of Medicine, Doctoral Programme in Clinical Research,
University of Helsinki
Helsinki, Finland

SMALL CHOROIDAL MELANOMAS

MANAGEMENT AND PROGNOSIS

Susanna Jouhi
nee Salkola

DOCTORAL DISSERTATION

To be presented, with the permission of the Faculty of Medicine,
University of Helsinki, for public examination in Lecture Hall 1 of
the Haartman Institute, Haartmaninkatu 3, on December 18, 2020, at 12 noon.

Supervised by

Professor Tero T. Kivelä

Department of Ophthalmology, University of Helsinki and Helsinki University Hospital,
Helsinki, Finland

Reviewed by

Professor Kai Kaarniranta

Department of Ophthalmology, University of Eastern Finland and Kuopio University
Hospital, Kuopio, Finland

Docent Ann Maria Paula Schalenbourg

Faculté de biologie et médecine, Hôpital ophtalmique Jules Gonin and L'Université de
Lausanne, Lausanne, Switzerland

Opponent

Professor Antonia Joussen

Department of Ophthalmology
Charité – Universitätsmedizin Berlin, Berlin, Germany

Cover image – Helsinki University Hospital, Department of Ophthalmology

The Faculty of Medicine uses the Urkund system (plagiarism recognition) to examine all
doctoral dissertations.

Dissertationes Scholae Doctoralis Ad Sanitatem Investigandam Universitatis Helsinkiensis

ISBN 978-951-51-6689-0 (pbk.)

ISBN 978-951-51-6690-6 (PDF)

ISSN 2342-3161 (pbk.)

ISSN 2342-317X (PDF)

Layout: Tinde Päivärinta/PSWFolders Oy/Ltd

Hansaprint, 2020, Vantaa, Finland

To Lauri, Kaarlo, and Okko

*“It is not the ship so much as the skilful sailing
that assures the prosperous voyage”*

George William Curtis

Table of contents

1. List of original publications.....	6
2. Abbreviations.....	7
3. Abstract.....	8
4. Summary in Finnish.....	10
5. Introduction.....	12
6. Review of the literature.....	14
6.1. Overview of choroidal melanoma	14
6.1.1. Epidemiology	14
6.1.2. Pathogenesis.....	14
6.1.3. Host- and environment-related characteristics.....	15
6.1.4. Growth patterns.....	16
6.1.5. Classification of small choroidal melanoma	16
6.2. Diagnosis	17
6.2.1. Diversity of small choroidal lesions	17
6.2.2. Clinical risk factors for choroidal melanoma	17
6.2.3. Clinical diagnosis	19
6.2.4. Observation for growth	21
6.2.5. Biopsy.....	21
6.3. Prognosis and prognostic factors	22
6.3.1. General aspects of prognosis	22
6.3.2. Histopathological prognosticators	22
6.3.3. Cytogenetic prognosticators	23
6.4. Treatment.....	24
6.4.1. Choice of primary treatment	24
6.4.2. Ruthenium ¹⁰⁶ brachytherapy.....	24
6.4.2.1. Radiation with ruthenium ¹⁰⁶	24
6.4.2.2. Radioactive applicators	25
6.4.2.3. Surgical technique.....	25
6.4.2.4. Local tumour recurrence	26
6.4.2.5. Ocular side-effects of radiotherapy	26
6.4.2.6. Vision outcome	27
6.4.3. Transpupillary thermotherapy.....	27
6.4.4. Proton beam therapy.....	28
6.5.5. Experimental treatments.....	29
6.6. Metastasis and survival	29
7. Aims of the study	31

8. Material and methods	32
8.1. Patients and study design	32
8.1.1. Studies I and II	32
8.1.2. Study III	32
8.1.3. Study IV	32
8.2. Primary treatment	33
8.2.1. Brachytherapy (Studies I and II)	33
8.2.2. Transpupillary thermotherapy (Study IV)	33
8.3. Data collection	34
8.3.1. Clinical evaluation	34
8.3.1.1. Studies I and II	34
8.3.1.2. Study III	35
8.3.1.3. Study IV	35
8.3.2. Calculations of growth rates and doubling times	36
8.3.3. Assessment of outcomes (Studies I and II)	37
8.3.4. Assessment of survival and metastatic status (Studies I and II)	37
8.4. Statistical methods and data analysis	37
9. Results	39
9.1. Treatment with the 10-mm or 15-mm plaque (Studies I and II)	39
9.1.1. Baseline patient, eye, and tumour characteristics	39
9.1.2. Plaque positioning	41
9.1.3. Local tumour recurrence and survival	41
9.1.4. Radiation maculopathy and optic neuropathy	42
9.1.5. Preservation of visual acuity	43
9.2. Small metastasizing choroidal melanomas (Study III)	43
9.2.1. Tumour Size and Characteristics	43
9.2.2. Local recurrence, development of metastases, and survival	45
9.3. Incipient choroidal melanomas (Study IV)	45
9.3.1. Baseline characteristics	45
9.3.2. Tumour characteristics at diagnosis	45
9.3.3. Transpupillary thermotherapy	47
10. Discussion	48
10.1. Treatment with the 10-mm or the 15-mm plaque	48
10.2. Small fatal choroidal melanomas	49
10.3. Diagnosis of incipient choroidal melanomas	50
10.4. Study strengths and limitations	51
10.5. Future aspects	51
11. Conclusion	53
12. Acknowledgements	54
13. References	56
Original publications	

1. List of original publications

This thesis is based on the following publications:

- I Salkola S, Heikkonen J, Eskelin S, Kivelä T,. Management of choroidal melanomas less than 10 mm in largest basal diameter with a 10-mm ruthenium¹⁰⁶ plaque. *Retina*. 2014; 34:2110-20.
- II Jouhi S, Reijonen V, Heikkonen J, Raivio V, Täll M, Kivelä T,. Brachytherapy of choroidal melanomas less than 10 mm in largest basal diameter: Comparison of 10-mm and 15-mm ruthenium¹⁰⁶ plaques. *Ophthalmology*. 2020; [Online ahead of print]
- III Jouhi S, Jager M, de Geus S, Desjardins L, Eide N, Grange J, Kiilgaard J, Seregard S, Midenä E, Parrozzani R, Caujolle J, Rospond-Kubiak I, Kivelä T,. The small fatal choroidal melanoma study: A survey by the European ophthalmic oncology group. *American Journal of Ophthalmology*. 2019; 202:100-8.
- IV Jouhi S, Al-Jamal R, Täll M, Eskelin S, Kivelä T,. Presumed incipient choroidal melanoma: Diagnostic criteria and management with transpupillary thermotherapy. [Submitted for publication]

The original publications are referred to in the text by their Roman numerals. They are reprinted in this thesis with the permission of the copyright holders.

2. Abbreviations

AJCC	American Joint Committee on Cancer
BAP1	BRCA-associated protein-1
BCVA	Best corrected visual acuity
BRCA	Breast cancer gene
CCA	15-mm round radioactive plaque
CCX	10-mm round radioactive plaque
CI	Confidence interval
CM	Choroidal melanoma
COMS	Collaborative Ocular Melanoma Study
CPI	Checkpoint inhibitor
CT	Computed tomography
DD	Disc diameter (about 1.5 mm)
DT	Doubling time
EDI	Enhanced depth imaging
FAF	Fundus autofluorescence
FAG	Fluorescein angiography
GEP	Gene expression profile
HPF	High power field
I ¹²⁵	Iodine ¹²⁵ isotope
ICGA	Indocyanine green angiography
IOP	Intraocular pressure
LBD	Largest basal diameter
LFT	Liver function test
MBD4	Methyl-CpG-binding domain 4
MRI	Magnetic resonance imaging
OCT	Optical coherence tomography
OOG	Ocular Oncology Group
OP	Orange pigment
Pd ¹⁰³	Palladium ¹⁰³ isotope
PRAME	Preferentially expressed antigen in melanoma
RPE	Retinal pigment epithelium
Ru ¹⁰⁶	Ruthenium ¹⁰⁶ isotope
SD	Standard deviation
SE	Standard error
SFCM	Small fatal choroidal melanoma
SRF	Subretinal fluid
SS	Swept source
TDT	Tumour doubling time
TNM	Tumour, Node, Metastasis classification
TTT	Transpupillary thermotherapy
UM	Uveal melanoma
US	Ultrasound
WHO	World Health Organization

3. Abstract

The first part of this thesis summarizes the current literature on small choroidal melanomas (CMs). Although the aetiology of this disease remains unknown, many risk factors have been reported. Differentiation from the more common choroidal naevi is generally clinical, and several typical characteristics of the malignant tumour have been used as a diagnostical tool. When the differentiation cannot be made clinically or by biopsy, the remaining option is to observe for growth that signals malignancy.¹⁻³ Small posterior tumours, in particular, have frequently been observed for growth because treatment would most likely affect vision. However, observation before treatment might increase the risk for metastases.⁴

Small CMs are typically treated conservatively, and enucleation is no longer a generally accepted alternative, partly because these tumours are thought to be able to micrometastasise years before diagnosis.⁵ The primary aims when managing small malignant CMs are to destroy the tumour and prevent local recurrences that might be associated with an increased risk for metastases.⁶⁻⁸ Eye-preserving treatments frequently allow for the preservation of useful vision as well. Several options are available for treating small CMs. In Finland, the majority of CMs are treated with episcleral plaque brachytherapy. Ruthenium¹⁰⁶ brachytherapy is an effective choice for small melanomas because the dose distribution is ideal for treating tumours with a thickness of up to 5.4 mm.⁹

The second and main part of this thesis summarizes the recent Finnish contributions to this field. The research presented in this thesis starts by retrospectively examining the treatment results from 10-mm ruthenium plaque (Study I), leading to a comparative study looking at the treatment results of both 10-mm round radioactive plaque (CCX) and 15-mm round radioactive plaque (CCA), with the aim to see whether and when it is safe to reduce vision-threatening radiation side-effects by using a smaller plaque (Study II) when treating small CMs less than 10 mm in their largest basal diameters (LBDs). The smallest 10-mm plaque delivers less scattered radiation compared with a 15-mm plaque or larger. Previously, it was not known whether a smaller radiation area and less scattered radiation could affect the recurrence rate, preservation of useful vision, and frequency of side-effects. This thesis found that the recurrence and complication rates were comparable between patients treated with the 10-mm or 15-mm plaque, but vision was more effectively preserved after treatment with the 10-mm plaque when the tumour was located close to the foveola. An eccentric location of the plaque was a risk factor for a local recurrence with both plaques. This study supports the treatment of small posterior CMs close to fovea with 10-mm rather than 15-mm plaque.

The challenge of diagnosing small CMs before they have the capacity to disseminate led to the search for the smallest size of a CM that had been reported to metastasize and an attempt to find out whether melanomas that disseminated and melanomas that did not

disseminate differed from each other clinically. As the Finnish population is not sufficiently large to collect data for this study, a retrospective collaboration study was done with the European Ocular Oncology Group (OOG) (Study III). Ten ocular oncology services submitted the data on all their patients with a CM of 3 mm or less in thickness and 9 mm or less in LBD who were treated and who subsequently developed metastases. This study found that CMs less than 3 mm in LBD are unlikely to metastasize. Observation without treatment beyond this limit might impair survival. Clinical characteristics of small fatal choroidal melanomas (SFCMs) did not differ from those of non-fatal CMs of a similar size: clinical characteristics predicting metastasis could not be identified.

Due to inspiration from Study III, ways to alternatively diagnose small melanomas before they get the capacity to disseminate were examined. The growth rates and tumour doubling times (TDT) of incipient CMs were calculated (Study IV). Short TDT and a fast growth rate combined with the calculated age at tumour origin after puberty supported a melanoma diagnosis. These were the nine smallest presumed CMs reported at that point. The earlier diagnosis enabled treating these patients with less invasive transpupillary thermotherapy (TTT), of which preliminary results are shared in this thesis. The reported growth parameters could be employed as diagnostic criteria for the incipient melanomas.

4. Summary in Finnish

Tämän väitöskirjan ensimmäinen osa kokoaa kirjallisuuskatsaukseksi suonikalvon melanoomaa käsittelevät ajankohtaiset julkaisut. Vaikka suonikalvon melanooman etiologia on avoin, monia sille altistavia tekijöitä tunnetaan. Melanooman erottaminen yleisemmästä hyvänlaatuisesta luomesta tapahtuu useimmiten kasvaimen kliinisten piirteiden perusteella. Jos kasvaimen pahanlaatuisuutta ei voida kliinisten piirteiden tai biopsian avulla varmentaa, kasvainta seurataan mahdollisen kasvun havaitsemiseksi.¹⁻³ Erityisesti pieniä silmän takaosassa sijaitsevia kasvaimia seurataan diagnoosin ollessa epävarma, sillä hoito todennäköisimmin heikentää näköä. Kasvainten seuraaminen ennen hoitoa voi suurentaa riskiä etäpesäkkeille.⁴

Pienet suonikalvon melanoomat hoidetaan tavallisesti paikallisesti eikä silmänpoisto ole enää yleisesti hyväksytty hoitokäytäntö, koska näiden syöpäkasvainten ajatellaan olevan leviämiskykyisiä jo vuosia ennen diagnoosia.⁵ Hoidon tärkein tavoite on tuhota kasvain ja estää kasvaimen paikallinen uusiutuma, mikä itsessään on riski etäpesäkkeille.⁶⁻⁸ Silmän säästävä hoito voi myös mahdollistaa käyttökelpoisen näkökyvyn säilyttämisen. Useista hoitovaihtoehdoista Suomessa on yleisimmin käytössä kovakalvon pinnalle asetettava sädehoitolevy. Rutenium¹⁰⁶-isotooppi soveltuu hyvin alle 5.4 mm paksujen kasvainten hoitoon ideaalisen annosjakaumansa vuoksi.⁹

Tämän väitöskirjan toinen osa kokoaa viimeisimmät suonikalvon melanoomien diagnostiikkaa ja hoitoa käsittelevät kotimaiset julkaisut yhdeksi kokonaisuudeksi. Tässä väitöskirjassa esitetty tutkimus alkoi hoitotuloksien selvityksellä, jossa pienten alle 10 mm halkaisijaltaan olevien kasvainten hoitoon oli käytetty 10-mm halkaisijaltaan olevaa ruthenium-sädehoitolevyä (osatyö I). Tutkimusta jatkettiin tekemällä vertaileva työ 10 mm (CCX) ja 15 mm sädehoitolevyillä (CCA) saaduista hoitotuloksista (osatyö II). Pienemmän halkaisijaltaan 10 mm levyn tavoitteena on säästää näkökykyä, koska pienemmän säteilevän alueen lisäksi pienempi levy tuottaa puolet vähemmän hajasäteilyä. Aiemmin ei ollut tiedossa vaikuttaako pienempi sädehoidettava alue ja pienempi hajasäteilyn määrä hoitotuloksiin niin kasvaimen uusiutumisen kuin näkökyvyn säilymisenkin suhteen. Paikallisissa kasvainten uusiutumisissa tai hoitoon liittyvissä komplikaatioissa ei havaittu eroa ryhmien välillä, mutta näkökykyä onnistuttiin säilyttämään hieman paremmin, kun hoitoon valittiin 10 mm levy. Tämä tutkimus suosittelee pienten silmän takaosassa sijaitsevien alle 10 mm halkaisijaltaan olevien kasvaimien hoitoon 10 mm halkaisijaltaan olevaa levyä.

Pienten suonikalvon melanoomien hoidon haasteena on diagnosoida pahanlaatuiset kasvaimet ennen kuin ne saavuttavat leviämiskyvyn. Asian selvittämiseksi käynnistettiin monikeskustutkimus, jossa 10 eurooppalaista silmäkasvaimia hoitavaa keskusta lähetti tiedot alle 3 mm paksuista ja alle 9 mm halkaisijaltaan olevista suonikalvon melanoomista,

jotka olivat lähettäneet etäpesäkkeitä. Aineistosta selvitettiin pienimmän etäpesäkkeitä lähettäneen suonikalvon melanooman koko ja etäpesäkkeitä lähettäneiden kasvainten kliinisiä piirteitä verrattiin kasvaimiin, jotka eivät olleet levinneet. Tulosten perusteella alle 3 mm halkaisijaltaan olevat kasvaimet eivät todennäköisesti vielä lähetä etäpesäkkeitä. Kasvavan muutoksen seuranta tämän raja-arvon jälkeen voi altistaa potilaan etäpesäkkeille. Kasvainten kliinisten piirteiden perusteella ei voi arvioida etäpesäkkeiden kehittymisen riskiä.

Selvitystä siitä, onko suonikalvon melanoomat mahdollisesta diagnosoida ennen kuin ne saavuttavat leviämiskyvyn jatkettiin. Joukko pieniä suonikalvon muutoksia, jotka ovat nopean kasvutaipumuksensa vuoksi todennäköisimmin melanoomia analysoitiin. Muutosten kasvunopeudet ja kahdentumisajat laskettiin ja kasvunopeuden perusteella arvioitiin potilaan ikä, jolloin kasvain olisi ollut havaittavissa. Nopeat kasvunopeudet ja kahdentumisajat verrattuna hyvänlaatuisiin luomiin ja laskennallinen ikä luomien havaitsemiselle aikuisiällä tukevat melanomadiagnoosia. Aiempaa varhaisempi diagnoosi mahdollistaa potilaiden hoitamisen transpupillaarisella lämpölaserhoidolla sädehoidon sijaan, jolloin voidaan välttyä säteilyn aiheuttamilta haittavaikutuksilta. Raportoitua tapaa mitata kasvunopeutta voidaan käyttää pienimpien suonikalvon melanoomien diagnostiikan apuna.

5. Introduction

Uveal melanoma (UM) is the second most common intraocular malignancy after retinoblastoma worldwide but the most common in Caucasians and adults.¹⁰⁻¹⁴ Up to one half of patients with UM develop metastatic disease within 10 years of diagnosis¹⁵⁻²¹, and the median survival after metastases has ranged from 6 to 13 months.^{19,22,23} Early diagnosis and treatment of UM are crucial because a larger size impairs prognosis.^{14,24-30} However, differentiation between benign naevi and small malignant CM can be difficult, as there is a spectrum of small choroidal melanocytic lesions ranging from benign lesions – naevi – with minor growth potential and no clinical risk factors to obviously malignant lesions with documented growth, several risk factors, and significant risk for metastatic disease.^{29,31,32} Management of small choroidal tumours is further complicated by the small number of naevi that transform into cancer.^{26,29,33}

Treatment of low-risk small choroidal melanocytic lesions is controversial because the significant majority of such tumours tend to remain stable and are likely benign choroidal naevi that will not spawn a melanoma during the lifetime of the patient.^{1,34-38} Certain clinical factors have been shown to predict growth that helps clinicians to distinguish small CMs from benign lesions.^{28,29,31,32,39-41} Such factors include, in particular, tumour thickness over 2 mm, subretinal fluid (SRF), symptoms, orange pigment (OP), and the margin either touching or being within 2 disc diameters (DD) of the optic disc.^{27,42} When risk factors are lacking, these tumours are followed-up in order to detect growth to ascertain malignancy and to justify treatment.³⁴ Particularly small posterior tumours are frequently observed for growth because treatment will most likely compromise vision. Tumour growth over a short period is considered a hallmark of CM although benign choroidal naevi can demonstrate slow growth over a period of many years to several decades as well, and the presence of such a slow growth in the absence of other risk factors is not an indicator for treatment.⁴³

Follow-up without treatment may theoretically have an effect on prognosis and expose the patient to metastatic disease if the tumour is a melanoma.⁴ The dilemma is that, while potentially malignant tumours that can spread systemically should be treated without delay, their identification can be challenging. Additionally, treatment of benign tumours does not benefit the patient and will likely deteriorate the patient's vision. The goal is to identify and only treat melanocytic lesions that are likely to spread to thus destroy these lesions before they do so.

A majority of CMs are treated conservatively either with episcleral plaque radiotherapy alone or in combination with TTT.⁴⁴⁻⁵⁰ Enucleation does not improve prognosis because tumours that have the capacity to disseminate are estimated to micrometastasise years before diagnosis.⁵ The primary goals in the management of small malignant melanomas are to destroy the tumour and to prevent local recurrences that might be associated

with an increased risk for metastases.^{6,51} Conservative treatment frequently allows for the preservation of useful vision. However, small CMs are frequently located posteriorly where the risk of vision loss is higher because of treatment side-effects. Tumours must be effectively treated to a sufficient degree to prevent local and systemic recurrence, simultaneously minimizing toxicity to healthy tissues.

In TTT, 810 nm diode laser energy is directed through a dilated pupil to the tumour apex, causing hyperthermia.⁴⁸ As a primary treatment TTT is controversial because of high local recurrence rates.⁵²⁻⁶⁰ It could be a treatment of choice in a selected group of patients with small choroidal tumours with minimal risk factors located close to but outside the central macula.^{53,61}

The research behind the present thesis aimed to determine the size limit of a small choroidal melanoma becoming capable of dissemination, describe the clinical tumour characteristics of metastasizing small tumours, and report TDTs and growth rates of incipient tumours in order to discover diagnostic criteria that could establish when small pigmented tumours should be treated. This thesis further reports the results of treatment with primary Ru¹⁰⁶ brachytherapy using 10-mm and 15-mm plaques and presents the comparison between the two plaque types to find the safest and most efficient treatment technique for small choroidal CMs.

6. Review of the literature

6.1. OVERVIEW OF CHOROIDAL MELANOMA

6.1.1. Epidemiology

The incidence of UM varies by race, latitude, age, and gender. It predominantly affects white Caucasians.⁶²⁻⁶⁴ The mean age-standardized incidence rate is 5.1 cases per million per year.¹⁵ The incidence rate varies inside Europe, with the highest being in Northern Europe at approximately 8–9.5 per million and, in Finland, as high as 12 per million⁶⁵, as compared to 4–6 million in Central Europe and 1.7–2 per million in Southern Europe.^{66,67} In the United States and Canada, the mean incidence rate is 4–6 per million,^{14,15,62,68,69} with the highest incidence rate occurring in non-Hispanic whites at 6 per million as compared to black Americans and Asians at 0.3 and 0.4 per million, respectively.⁷⁰ The incidence rate in Australia is as high as in Northern Europe at 8 per million.⁷¹ Standardized incidence rate estimates suggest that 7,000 new patients are diagnosed annually worldwide.⁶⁶ The standardized incidence of UM has remained stable over the years^{14,15,20,72,73} in contrast with the rising incidence of cutaneous⁷⁴ and conjunctival melanoma⁷⁵. However, the incidence of UM in Finland has increased in 20 years from approximately 8 to 12 per million due to an increasing overall population age.^{65,67} The incidence rate increases with age, peaks at the age of 70¹⁰⁻¹², and plateaus after 75 years of age^{12,14,15,66}. The median age range at diagnosis is 59–62 years.^{12,15,64,76} The median age slowly increased over the years to 62 years due to increasing life expectancy.⁷⁷ The incidence of UM in children and teenagers is rare⁷⁸⁻⁸⁰ and increases steadily until the age of 11 years, followed by a transition to a more than 10-times faster increase after the age of 17.⁸¹ Males show a slight predominance in crude incidence.^{10,64,67}

6.1.2. Pathogenesis

The aetiology of UM is unknown.⁶³ However, five driver mutations have been commonly found: BAP1, GNAQ, GNA11, SF3B1, and EIF1AX.⁸²⁻⁸⁹ UMs originate from neuroectodermal melanocytes in the middle, uveal layer of the eye wall.⁹⁰ UMs are divided into different types according to their location: anteriorly located iris melanomas and posteriorly located ciliary body melanomas and CMs. The majority of UMs originate in the choroid (85%–90%), and remainder originate in the iris (3%–5%) and the ciliary body (5%–8%).^{67,76,91,92}

UMs, although also originating from melanocytes, differ from cutaneous melanoma and mucous membrane – including conjunctival – melanomas in terms of epidemiology, aetiology, biology, genetics, and clinical features, including a high propensity to metastasize haematogenously to the liver.⁹³ UM predominantly disseminates haematogenously because there is no traditional lymphatic drainage within the eye.⁹⁴

6.1.3. Host- and environment-related characteristics

Several predisposing factors have been identified as affecting the occurrence of UM, including, as aforementioned, race, age, and gender and additionally including blue eyes, fair skin, the inability to tan, ocular or oculodermal melanocytosis, choroidal naevi, and germline mutations in BAP1 or MBD4 genes.

Although light skin and a blue iris colour are established risk factors for UM⁹⁵, the role of sun exposure is weak,^{63,96,97} in contrast to cutaneous melanoma.⁹⁸ Long-term exposure to ultraviolet-radiation is not associated with the incidence of UM,⁹⁹ whereas intermittent exposure to ultraviolet rays from welding has been reported to double the risk of the development of UM,^{99,100} this finding is, however, questionable.¹⁰¹

Congenital ocular melanocytosis is an important condition predisposing individuals to UM.¹⁰² It is a rare developmental pigmentary disorder in which the number of melanocytes is increased in ocular and periocular tissues. Melanomas arising from ocular melanocytosis are associated with more frequent genomic alterations and a double risk for metastatic disease.^{103,104}

Choroidal naevi carry a small risk of transformation into CM.^{27-29,31,32,39-41,105-107} At least one in 10 melanomas originates from pre-existing naevi,¹⁰⁸⁻¹¹⁰ which, however, exist in approximately 3–8% of the general population.^{25,26,111,112} The annual rate for malignant transformation is estimated at one in 3,664 to 8,845,^{26,113} and the lifetime risk for malignant transformation is about 1%.¹⁰⁷

Although UM generally occurs sporadically,⁶³ BAP1 germline mutations have been described in families with hereditary BAP1 cancer syndrome, including UM, comprising 0.6–6 % of UM patients depending on age.¹¹⁴⁻¹²⁴ The prevalence of pathogenic BAP1 variants is reported to be 25% in Finnish familial UMs.¹²⁵ Patients with BAP1 cancer syndrome have an increased risk of several cancers, the most frequent of which is UM.^{82,115,117} Patients with UM have over 10% risk for other malignancies, which is, to a certain extent, driven by the presence of germline BAP1 mutations.¹²⁶

UMs are associated with a predisposing germline loss-of-function variant in the MBD4 gene causing hypermutated tumours¹²⁷⁻¹²⁹, a variant which has a prevalence in UM patients that is 9.15 times more frequent than in the general population.¹²⁷ These hypermutated tumours should be recognized since they may respond to checkpoint inhibitor (CPI) immunotherapy.^{118,130-132}

6.1.4. Growth patterns

Small choroidal tumours are frequently flat or crescent in shape. After growing vertically, they become dome-shaped, and when the Bruch's membrane ruptures, they extend into the subretinal space and become mushroom-shaped.^{90,133} The sclera presents resistance to the tumour expansion, and the tumour therefore protrudes into the vitreous cavity.¹³⁴ CMs may eventually extend through existing scleral channels along nerves and vessels into the episclera or the orbit, an event which occasionally occur, even with small choroidal tumours around the optic disc,¹³⁵⁻¹³⁹ an area that is rich with intrascleral nerves and vascular channels.^{134,140,141}

6.1.5. Classification of small choroidal melanoma

Numerous definitions have been devised for each CM size category.^{18,142} The use of classifications should allow for the comparison of treatments for equivalently sized and staged tumours. The Collaborative Ocular Melanoma Study (COMS) has defined a small CM as being 1–2.5 mm in thickness and 5–16 mm in LBD; tumours smaller than this are regarded as probable naevi.¹⁴³ It has been estimated that when the LBD is 5–6 mm, about 70 choroidal naevi are diagnosed for each melanoma, and it has been stated that few melanomas would be less than 5 mm in diameter.^{33,144}

UMs have been classified by size and anatomic extent which affect the risk for metastases and survival.²⁴ According to the American Joint Committee on Cancer (AJCC) Tumour, Node, Metastasis (TNM) staging systems latest 7th and 8th edition,^{145,146} the criteria of tumour size category T1 are a thickness less than 3.0 mm if the LBD is less than 12.0 mm or a thickness less than 6.0 mm if the LBD is less than 9.0 mm. Tumour size category T2 tumours can be classified as medium-sized, and the corresponding size criteria are a thickness less than 3 mm if the LBD is 12.1–18 mm, a thickness of 3.1–6 mm if the LBD is 9.1–15 mm or a thickness 6.1–9 mm if the LBD is 9.1–12 mm.¹⁴⁵

Table 1. Current Tumour, Node, Metastasis categorization of choroidal and ciliary body melanomas according to anatomical extent by the American Joint Committee on Cancer¹⁴⁵

T1	Tumour Size Category 1
T1a	T1 category without ciliary body involvement or extraocular extension
T1b	T1 category with ciliary body involvement
T1c	T1 category without ciliary body but with extraocular extension 5 mm
T1d	T1 category with ciliary body and extraocular extension 5 mm
T2	Tumour Size Category 2
T2a-d	T2 category and similar sub-categories as for T1
T3	Tumour Size Category 3
T3a-d	T3 category and similar sub-categories as for T1
T4	Tumour Size Category 4
T4a-d	T4 category and similar sub-categories as for T1
T4e	Any tumour size category with extraocular extension > 5 mm

6.2. DIAGNOSIS

6.2.1. Diversity of small choroidal lesions

The difficulty of detecting small CM relates to the diversity of small pigmented choroidal lesions ranging from benign naevi, via indeterminate pigmented lesions, to malignant melanoma.^{28,31,32,147} Clinical risk factors help clinicians to determine which small melanocytic choroidal tumours should be treated or biopsied without waiting for tumour growth. Lesions carrying any of these risk factors are generally referred for evaluation by a retinal specialist or ocular oncologist. Smaller melanomas typically show fewer risk factors than larger melanomas, making their diagnosis more difficult.¹³³

There is no consensus on indications for treatment of small choroidal melanocytic tumours. The more risk factors are present, the more likely it is that a choroidal lesion is a melanoma.¹⁴⁸ Lesions that display one risk factor have a 38% risk for growth, and those with two or more factors show growth in over 50% of tumours.¹⁴⁸ When a lesion does not have any of these factors, the risk for growth and malignancy over 5 years is approximately 3%, and the tumour is more likely to be a choroidal naevus that can be monitored periodically.¹⁴⁸ However, the typical risk factors^{27,42} are not pathognomonic for malignancy.¹⁴⁹ The treatment of low-risk small choroidal lesions is controversial because a majority of these lesions will remain stable and likely be benign choroidal naevi.^{2,28,31,53}

6.2.2. Clinical risk factors for choroidal melanoma

Certain clinical signs are risk factors for tumour growth and metastasis.^{27,31,32,39-42,150} These independent risk factors are a tumour thickness over 2 mm, a LBD over 5 mm, SRF, ocular symptoms, OP, a posterior tumour margin either touching or located within 3 mm of the optic disc, ultrasound (US) hollowness, the absence of a halo around the tumour, and the absence of drusen (Fig. 1).

A tumour thickness over 2 mm (Fig. 1) is a consistent independent risk factor for malignancy^{27-29,33,150} and is associated with the class 2 gene expression profile (GEP)¹⁴⁹ (see below – 6.3.3: Cytogenetic prognosticators). Tumours less than 1.0 mm in thickness are most likely naevi²⁷ and there are approximately 125 naevi for each melanoma in the thickness range between 1.5–2 mm, but only 25 naevi for each melanoma in the range of 2–2.5 mm, and 5 in the range of 2.5–3 mm.¹⁴⁴ The rate of melanoma-related metastasis at 10 years gradually increases by each millimetre from 6% to 51% when thickness increases from the 0–1.0mm to >10.0mm, respectively.⁷⁶ Additionally, a tumour diameter over 5 mm has been described as an independent risk factor for malignancy.¹⁵⁰

Clinically detectable SRF (Fig. 1) is a strong indicator of metabolic activity and malignancy.^{27,151} Optical coherence tomography (OCT) evidence of SRF has a predictive value in identifying tumours that have cellular activity and are likely to grow. However,

around 10% of choroidal naevi have been reported to present small amounts of SRF detectable on OCT due to the gradual atrophy of the retinal pigment epithelium (RPE) and loss of its pumping action that normally keeps the subretinal space dry.^{64,150,152-155}

Small choroidal tumours can cause visual symptoms when located close to or extending under the fovea (Fig. 1). Since UMs are generally slowly growing tumours,^{20,156} these symptoms are generally not specific. The most commonly presenting symptoms are blurred vision, photopsia, floaters, a visual field defect, and metamorphopsia.^{27,108} The development of symptoms is frequently indicative of growth and frequently leads to the diagnosis of melanoma.¹⁵⁷

The RPE can display intracellular lipofuscin accumulation overlying active tumours (Fig. 1).^{158,159} This is clinically evident with biomicroscopy or fundus autofluorescence (FAF) of the posterior pole and is described as OP over the pigmented lesions. OP appears black over amelanotic tumours. OP can be found in over 6–10% of benign posteriorly located naevi, which should be frequently evaluated for malignancy; thus, the presence of OP alone does not automatically indicate malignancy.¹⁰⁵

A posterior tumour margin either touching or extending within 2 DD from the optic disc margin has been associated with a higher risk for malignancy.^{27,42} However, this location was reportedly no longer a significant risk factor when the analysis was based on multimodal imaging.³³

Small CMs typically have a relatively low acoustic reflectivity on A- and B-scan US due to their homogenous regular structure (Fig. 1).¹⁶⁰⁻¹⁶² Additionally, a regular acoustic internal structure and the presence of a vascularity within the lesion can be detected.^{160,163} These are useful when differentiating between choroidal metastases and haemangiomas.¹⁶⁴ B-scan US is additionally instrumental in measuring tumour thickness and the LBD, as well as in exploring tumour shapes and the presence of an extrascleral extensions.^{133,165} Extraocular extensions of CM, which are rare in small melanomas,¹⁶⁶ appear as echolucent nodules adjacent to the sclera at the tumour base.¹³³ Choroidal naevi are frequently excessively thin to be reliably evaluated with US and do not facilitate the diagnosis of thin melanomas less than 1 mm in thickness either.¹³³

The absence of a halo or overlying drusen are risk factors for growth.⁴² A halo is a ring of lighter pigmentation surrounding a choroidal lesion. This phenomenon is generally associated with naevi but can occasionally be seen with melanomas.¹⁶⁷ Drusen are an age-related sign of a compromised RPE that have developed over a long period of time. Drusens can be seen overlying choroidal lesions and are the signal of inactive, and therefore most likely benign, naevi. Drusen do not exclude the possibility of malignancy. They may occasionally suggest that a previous naevus has become active, particularly when they are unevenly distributed over the lesion.¹⁰⁵ Drusen can additionally be an incidental finding

over a choroidal tumour when they are further present in the surrounding retina as an age-related degeneration.




Figure 1. A small choroidal melanocytic tumour with 4 clinical risk factors: symptoms (metamorphopsia and a visual field defect), on ophthalmoscopy (**Left**): subretinal fluid (SRF) (white arrows), and orange pigment (OP) (black arrows), and on B-scan ultrasonography (**Right**): a tumour thickness over 2 mm, as well as a relatively hollow internal reflectivity.

6.2.3. Clinical diagnosis

Clinical risk factors are most frequently evaluated by comprehensive ophthalmic examination, US, OCT, and in the case of indeterminate tumours, by serial observations for tumour growth as well. Other available methods for diagnostic verification are different modalities of OCT, fluorescein angiography (FAG), indocyanine green angiography (ICG), and FAF. For small tumours, computed tomography (CT) or magnetic resonance imaging (MRI) is generally not needed unless an extrascleral extension or optic nerve invasion are suspected.

OCT and optical coherence tomography angiography (OCTA) are non-invasive imaging modalities that provide a more detailed image of the retina and choroid when compared to US.¹⁶⁸ OCTA provides insight into retinal vascular abnormalities related to CMs, such as an enlargement of the deep foveal avascular zone and a decrease in the superficial and deep parafoveal capillary vascular densities, which appear to be correlated with the presence of SRF and to be absent in choroidal naevi.^{168,169} OCT technology is traditionally limited to the examination of posterior lesions, but newer technologies enable a view of peripheral lesions as well.¹⁷⁰ Furthermore, a microscope-integrated OCT has been used intra-operatively to guide a biopsy in real-time, increasing the chance of obtaining an adequate sample and avoiding unnecessary damage.¹⁷¹

Enhanced depth imaging optical coherence tomography (EDI-OCT), uses spectral domain technology, and swept-source optical coherence tomography (SS-OCT) with faster

scanning engines and can image structures from the choroid to the inner sclera.¹⁷² SS-OCT uses longer wavelengths compared with earlier generation OCTs, enabling improved penetration through the RPE and examination of deeper structures.^{168,172}

EDI-OCT, as a result of a modification of the SD-OCT technique, enables the reliable imaging of the choroid in its full-thickness.^{173,174} By permitting high-resolution visualization of the choroid, EDI-OCT is sensitive to recognizing thin lesions and possibly detecting a submillimeter early melanoma that might not be apparent with US.¹⁷⁵⁻¹⁷⁷ Typical characteristics visible on EDI-OCT include deep optical shadowing, thinning or compaction of the choriocapillaris, disruption of the adjacent retinal photoreceptor layer, SRF with or without lipofuscin deposition, and intraretinal fluid.¹⁷⁷

FAF is a non-invasive technique useful for identifying lipofuscin in pigmented choroidal lesions.¹⁷⁸⁻¹⁸⁰ It is based on the stimulated emission of light from naturally occurring fluorophores, the most significant being lipofuscin.¹⁸⁰ A majority of naevi are not hyper-autofluorescent.¹⁸⁰ Naevi generally show a normal pattern of background FAF, but a minority of cases may reveal areas of a decreased FAF signal that is associated with chronic RPE degenerative features and atrophy.^{180,181} Nearly 90% of CMs located at the posterior pole show at least one focus of increased autofluorescence.¹⁸² SRF represents a barrier between the photoreceptors and the RPE, preventing their normal phagocytosis. As a result, they accumulate on the outer retinal surface and in the subretinal space and become a source of autofluorescence.¹⁸⁰

Invasive methods, such as FAG and ICGA, which involve the injection of an intravenous dye, have been used to differentiate melanoma from a masquerading pathology to search for secondary neovascularization or ischaemia or to assist in visualizing lesions obscured by media opacities, but non-invasive methods have replaced these methods to a certain extent.^{183,184} The imaging of the choroidal vasculature with ICGA may reveal microvascular patterns that can help to differentiate a naevus from a melanoma through the recognition of the typical features of melanomas, such as an irregularity of the lesion's margins; a heterogeneous, hyporeflective choriocapillaris plexus with avascular areas; hyperreflective choriocapillaris rings; thick choroidal vascular networks; and choroidal vascular loops.^{169,185} The presence of haemorrhages and lipid exudation on ICGA as well as FAG are useful for the differential diagnosis of a malignant CM and choroidal metastasis as opposed to peripheral exudative haemorrhagic chorioretinopathy.¹⁸⁶ Typical findings of CM on FAG include hypofluorescence with mottled hyperfluorescence in the arterial or early venous phase. However, a small CM with an intact overlying RPE can demonstrate no appreciable abnormalities.¹⁸⁷⁻¹⁸⁹ FAG can reveal OP clumps on the surface of the tumour, where these clumps appear to be hypofluorescent. SRF can be seen by the accumulation of fluorescein in the late frames, leading to hyperfluorescence.¹⁸⁷⁻¹⁸⁹

6.2.4. Observation for growth

Diagnosing a small CM may include observation with watchful waiting in the event that the diagnosis is not otherwise ensured.¹⁻³ Observation is thought to be justified and unable to impact prognosis³⁹ because it has been estimated that development of micrometastases initiates several years before the diagnosis of a CM.¹⁹⁰⁻¹⁹² However, it is not known when the dissemination takes place, but tumours as small as 3 mm in LBD and 1.5 mm in thickness have been predicted to be capable of dissemination.¹⁹³ Initial observation for growth might be appropriate in a majority of choroidal tumours less than 2.25 mm in thickness and in patients under 60 years of age since a majority of those tumors will belong to GEP class 1 (see below: 6.3.3: Cytogenetic prognosticators) and have a lower risk of metastases.¹⁴⁹

Not all enlargement, however, represents true transformation into melanoma. Slow enlargement over years at a median growth rate of 0.06 mm/year has been shown to likely represent a benign naevus without transformation over time.^{43,194} It is current practice to consider an enlargement over 0.5 mm within 2 years to be suggestive of a malignant tumour.¹⁹⁵

De novo tumours become visible after 20 cell divisions still being less than 1 mm in size.¹⁹³ The development of a new naevus in adulthood or asymmetric growth is generally cause for suspicion and should be considered as a *de novo* melanoma until proven otherwise.¹⁹⁶⁻¹⁹⁸

6.2.5. Biopsy

Small lesions without sufficient clinical evidence for malignancy are either observed for growth and, particularly, growth rates or, if sufficiently conspicuous, are biopsied before being defined and treated as melanomas.^{199,200} The biopsies today are performed more frequently for prognostic rather than diagnostic purposes.²⁰¹⁻²⁰⁵ Routine use of cyto- or histologic confirmation for diagnosis is not a current practice, unlike in a majority of other cancers, because of an over 95% accuracy rate of non-invasive diagnosis, at least for tumours thicker than 3.0–4.0 mm,^{17,108,161,206,207} whereas, as high as one-third of small CMs are initially misdiagnosed clinically.²⁰⁸ Despite the claimed risk for tumour dissemination secondary to diagnostic invasive approaches,²⁰⁰ a biopsy has been shown to be sufficiently safe and successful for tumours as thin as 0.7–1.0 mm²⁰⁹⁻²¹¹ and 2–3 mm^{149,209,212-215} in LBD.

However, a fine-needle aspiration or vitreous cutter biopsy does not exclude the possibility of a malignant tumour, including with a non-malignant result, because the sample might not be representative of the entire tumour, particularly of those that are transformed naevi.^{208,216,217} Conversely, a prognostic biopsy may occasionally return a wrong result.^{218,219}

6.3. PROGNOSIS AND PROGNOSTIC FACTORS

6.3.1. General aspects of prognosis

In the TNM classification, the anatomical extent of the primary tumour is marked with a T category. Its regional lymph node involvement, which is extremely rare with UMs,²²⁰⁻²²² is represented under the N, and eventual metastases with the M. Patients with evident regional lymph node or systemic metastases, irrespective of the tumour size, are staged as a stage IV. Ten-year Kaplan-Meier survival estimates for small CMs without ciliary body involvement or an extraocular extension (T1, stage I) range between 88–94% (95% CI, 84–96%) and around 80% (95% CI, 75–84%) for medium sized (T2, stage IIA) CMs.^{24,223}

Table 2. Current TNM stages of choroidal and ciliary body melanomas according to American Joint Committee on Cancer (AJCC)

Stage I	T1a	N0	M0
Stage IIA	T1b-d	N0	M0
	T2a	N0	M0
Stage IIB	T2b	N0	M0
	T3a	N0	M0
Stage IIIA	T2c-d	N0	M0
	T3b-c	N0	M0
	T4a	N0	M0
Stage IIIB	T3d	N0	M0
	T4b-c	N0	M0
Stage IIIC	T4d-e	N0	M0
Stage IV	Any T	N1	M0
	Any T	Any N	M1a-c

6.3.2. Histopathological prognosticators

Three main cell categories are microscopically distinguished in UM: spindle cell, epithelioid cell, and mixed.²²⁴ Spindle cell melanomas are associated with a better prognosis, mixed cell melanomas are associated with an intermediate prognosis, while epithelioid tumours grow faster, are more likely to metastasize, and are correlated with a shorter survival time.^{140,225-229} The 15-year mortality rates of patients with spindle, mixed, or epithelioid cell melanoma are 19–26%, 59–60%, and 72–75%, respectively.^{230,231} An increase in the number of mitoses per 40 HPF, a high fraction of PC-10 and Ki-67 indicating high cellular activity, a large mean diameter of the 10 largest nucleoli,²³²⁻²³⁴ a high microvascular density,^{232,235} an increased infiltration by lymphocytes²³⁶ and macrophages^{237,238}, high expression of insulin-like growth factor-1 receptor,²³⁹ a high expression of human leukocyte antigen I and II²⁴⁰, and the presence of extravascular matrix network patterns loops and networks²⁴¹⁻²⁴³ have been associated with a worse prognosis.^{224,225,242,244-247}

6.3.3. Cytogenetic prognosticators

The better prognosis of patients with small CMs compared to those with large CMs is likely related to fewer and less risky cytogenetic abnormalities of small CMs compared to larger tumors and, hence, a lower metastatic capacity.^{4,103} UM carries distinctive molecular pathway defects, with its own chromosomal and molecular alterations.²⁴⁸⁻²⁵⁰ Prognostic biopsies of conservatively treated UMs that allow for the analysis of their cytogenetic, gene expression, and molecular genetic features are not yet a routine part of the treatment of all UM patients, but are increasingly more common.

A complete or partial loss of chromosome 3, which occurs in approximately 50% of UMs, is associated with a reduction of 50% in survival.²⁵¹⁻²⁵⁶ This loss of chromosome 3 is generally associated with a multiplication of chromosome 8, 8q, or parts of 8q, and additional copies of chromosome 8q are correlated with a high mortality rate.^{251,252} Monosomy 3 together with gains in chromosome 8q is thus associated with the highest risk for metastasis, whereas either monosomy 3 or gains in chromosome 8q are associated with an intermediate risk and disomy 3 with the lowest risk.^{214,252,257,258} Chromosome 1 and chromosome 6 changes are frequent as well.²⁵⁹⁻²⁶¹ Loss of chromosome 1p correlates with a reduced survival rate,^{262,263} whereas a chromosome 6p gain is suggestive of a protective effect and associated with a better prognosis.²⁶⁴ The determination of the chromosomal status may be affected by prior irradiation.²⁶⁵

Several somatic mutations have been identified as having an influence on an UM patient's prognosis. Tumorigenesis is driven by mutually exclusive gain-of-function mutations in members of the Gq signalling pathway (GNAQ, GNA11, PLCB4, or CYSLTR2) followed by near-mutually exclusive mutations in BAP1, EIF1AX, SF3B1, or SRSF2.²⁶⁶⁻²⁶⁸

GNAQ and GNA11 mutations, members of the q class of the G-protein α -subunits, are found in all stages of UM, but additionally found in naevi.^{85,88} Three times as many metastatic UMs carry mutations in GNA11 as compared to GNAQ, suggesting that cells carrying a GNA11 mutation are designated for the metastatic process.^{83,84,269,270} Furthermore, PLCB4 and CYSLTR2 oncogenes are, similarly, a gain-of-function mutations leading to an activation of the same signalling pathway and thus promoting UM tumorigenesis.^{271,272}

BAP1 has been mapped on a tumour suppressor gene located on chromosome 3, and its loss is a strong indicator for worse prognosis.^{82,115,269,273} Whole-exome sequencing of metastatic UMs has identified inactivating somatic mutations in BAP1 in 84% of metastasizing tumours, suggesting that the inactivation of BAP1 is a pivotal event in the development of metastases.⁸² Mutations of the X-linked translation initiation factor EIF1AX have a protective effect and are associated with an 8-fold decreased risk of metastasis.^{87,269} Similarly, mutations in the splicing factor SF3B1 are correlated with improved survival

rates as well,^{86,87,274} although there have additionally been studies that have concluded the opposite argument for this association.^{87,269}

GEP is a technique which uses the mRNA of tumour biopsies in order to predict metastatic risk.^{254,275} Based on the results of their GEPs, UMs are divided into two major prognostic subgroups: those in the class 1 with a low (class 1A) or intermediate metastatic risk (class 1B) and those in class 2 with a high risk.^{114,203,254,276} Class 1 tumours are associated with disomy 3, and the gain of 6p, and class 2 tumours are associated with monosomy 3 and mutations in the BAP1.^{82-85,88,114,271,272} Metastatic rates have been reported to be as low as 1% in class 1 and up to 26% in class 2 cases at a median follow-up time of 17 months.²⁰³ Patients with a class 2 UM tend to be older, have thicker tumours with epithelioid cells as well as a high mitotic rates, and present BAP1 mutations.²⁰³ Although the majority of metastases occur in patients with class 2 tumours, class 1 tumours may give rise to metastasis as well.²⁰³ Preferentially expressed antigen in melanoma (PRAME) is an independent prognostic biomarker in UM that identifies increased metastatic risk in patients with class 1 or disomy 3 tumours.²⁷⁷

6.4. TREATMENT

6.4.1. Choice of primary treatment

A majority of CMs are conservatively managed with radiotherapy, although selected large tumours continue to be treated with enucleation^{8,278,279} and certain tumours undergo local resection^{280,281}. Conservative treatment of CM aims to destroy the tumour while preserving as much useful vision as possible.

Three major radiotherapeutic techniques are available: brachytherapy with various isotopes and radioactive plaques, external beam radiotherapy with photons from linear accelerators or gamma knives, and charged particle beams.²⁸²⁻²⁸⁹ All of these techniques are effective for CMs of all sizes and have high rates of local control with survival rates that are similar to those observed after enucleation.^{72,290,291} TTT can be a safe method for selected patients with thin and small tumours,^{52,53} but high recurrence rates with thicker tumours have been a concern.^{53,292}

6.4.2. Ruthenium¹⁰⁶ brachytherapy

6.4.2.1. Radiation with ruthenium¹⁰⁶

Treatment with a beta emitter Ru¹⁰⁶ applicator was first introduced in 1964 in former East Germany.^{45,293,294} It gradually became more popular, particularly within Europe in the 1970s to 1980s.²⁹⁵⁻³⁰² Ruthenium¹⁰⁶, with a half-life of 373.59 days, is suitable for CMs of up to 5.4 mm in thickness because the beta radiation has a limited depth of penetration.^{9,303} The advantage of Ru¹⁰⁶ over gamma emitters and proton beam when treating small CMs is

thought to be fewer radiation-related complications and thus better preservation of vision since it causes less damage to healthy tissue.^{287,304,305}

6.4.2.2. Radioactive applicators

Round episcleral plaques are bowl-shaped and 10–25 mm in diameter (Fig. 2).⁹ Notched plaques are available for juxtapapillary tumours and ciliary body tumours. Ruthenium¹⁰⁶ plaques are commercially manufactured as pure silver plaques with encapsulated radioactive Ru¹⁰⁶. Their outer surface is lined by a thick silver layer that absorbs more than 99% of the radiation to prevent irradiation of tissues surrounding the eye,³⁰⁶ whereas the silver layer of the active inner side has a thickness of 0.1 mm. Eyelets allow the plaque to be sutured to the sclera. The activity of the Ru¹⁰⁶ plaque depends on the size and shape of the plaque and ranges between 4.1–40.3 MBq, equal to 0.11–1.09mCi.⁹




Figure 2. Round radiation plaque ⁹

The radiation dose is delivered to the tumour during a precalculated and continuous time period. The planned dose and treatment period are directly proportional to the tumour thickness. The prescription point is determined by the tumour thickness, to which 1 mm is added for the sclera.

The difference between the doses delivered to the sclera and apex increases with increasing tumour height. The prescription dose to the apex is generally 80–120 Gy, and frequently aims to deliver at least 250 Gy to the sclera³⁰⁷, although certain centres prefer a scleral dose of 350Gy.³⁰⁸ Scleral doses as high as 1,500 Gy using Ru¹⁰⁶ have been administered without scleral necrosis,³⁰⁹ which, however, remains a possible complication with such high doses.

The conventional practice is to position the plaque centred over the tumour to ensure that it overlaps the tumour margins by at least 2 mm in all directions.^{6,282} The notion of scattered irradiation has induced certain centres to use eccentric plaque placement, with the posterior edge of the plaque aligned with the posterior tumour margin and without a safety margin in an attempt to save the fovea and optic nerve.^{307,308,310} A wide radiation safety margin inevitably increases side-effects, particularly when the posterior tumour margin extends close to the optic disc or fovea.

6.4.2.3. Surgical technique

Total or partial peritomy is performed to expose the underlying sclera, followed by a disinsertion of the muscles if the muscles are located in the tumour area in order to enable plaque positioning over the sclera. The tumour margins are identified, for example, by transilluminations or indirect ophthalmoscopy with indentation and are marked with diathermy or ink. For plaque positioning, a non-irradiating dummy plaque is used. Once it is placed correctly over the tumour, it is replaced with the radioactive plaque. Intra- and post-operative US can be useful to confirm the exact plaque positioning.^{311,312} A mattress

suture over the plaque is recommended but is impossible when treating posterior tumours with the smallest plaques. The plaque cannot be placed closer than approximately 1.5 mm from the margin of the optic disc because of the optic nerve sheath, which has a diameter of 5.1 mm behind the eye.³¹³ When the radioactive plaque is fixed to the sclera, any disinserted rectus muscles are resutured over the plaque, and the conjunctiva is closed. When the prescribed radiation dose has been delivered, the plaque is removed during a second procedure. Any disinserted rectus muscles are replaced in their anatomical position. The inferior oblique muscle is generally left unattached since it has frequently been only partially disinserted.

6.4.2.4. Local tumour recurrence

Ru¹⁰⁶ brachytherapy provides 5-year local recurrence rates from 2–15% for small and medium sized CMs when not restricted to a particular plaque size.^{303,307,314–317} A larger tumour size and posterior location are risk factors for recurrence.^{303,318} Although a majority of recurrences occur within the first few post-operative years, regrowth, including after 15 years, has been reported.³⁰⁹ An increase in tumour size after initial tumour regression may occasionally further result from an intratumoural haemorrhage and does not necessarily indicate a local recurrence.³¹⁹

6.4.2.5. Ocular side-effects of radiotherapy

CMs are resistant to radiation, and their treatment, therefore, requires high radiation doses that may result in radiation-related complications and impaired best corrected visual acuity (BCVA). Brachytherapy of small posterior CMs with Ru¹⁰⁶ leads to posterior segment complications such as radiation maculopathy, retinopathy and optic neuropathy, as well as vitreous haemorrhage, whereas anterior segment complications such as cataract and iris neovascularization, with or without neovascular glaucoma, are rare.^{308,309,317,320–323}

Radiation maculopathy appears clinically as retinal vascular changes such as microaneurysms, telangiectasias, retinal haemorrhages, microinfarcts, macular oedema, neovascularization, and lipid exudation.^{31,324,325} Diabetic patients are at a higher risk.³²⁶ Bevacizumab and intravitreal or periocular triamcinolone can transiently reduce macular oedema to maintain visual acuity (VA).^{311,327} Maculopathy caused by exudates and oedema can be treated by administering TTT to the tumour if the tumor is located outside the macula. The 5-year cumulative incidence rate of developing radiation maculopathy following Ru¹⁰⁶ brachytherapy is 30–55%.^{302,308,310,328}

Radiation optic neuropathy is a less common complication after treatment of small posterior tumours with Ru¹⁰⁶, presenting in up to 12% of cases after 5 years.³⁰² This risk is related to the distance of the posterior tumour margin to the optic disc and is minimal beyond 4 mm.^{328,329}

6.4.2.6. Vision outcome

Radiation retinopathy and maculopathy continue to be vision-limiting complications following radiation exposure.³³⁰ Ru¹⁰⁶ brachytherapy may allow to for the retaining of useful vision for a considerable period of time.³⁰⁰ Risk factors for losing vision after brachytherapy are older age, a lower initial BCVA, a tumour location close to the fovea or optic disc, a larger tumour thickness, and a temporal tumour location.^{307,309} Three to 5 years after brachytherapy, approximately 50% of patients maintain a BCVA of 20/200 (decimal scale, 0.1) or better, and 30% a BCVA of 20/50 (decimal scale, 0.4) or better in the tumour eye.^{309,331} When using eccentric placement, up to 75 % of patients with their posterior tumour edge at least 3 mm from the fovea may retain 20/40 (decimal scale, 0.5) BCVA at 4 years after brachytherapy.^{308,310} Vision can further improve after radiation due to a resolution of SRF following tumour regression.^{57-59,332,333}

When compared with Pd¹⁰³, Ru¹⁰⁶ brachytherapy is associated with improved preservation of vision.²⁸⁷ Following I¹²⁵ brachytherapy, a good local tumour control of approximately 90% at 10-years has been achieved for small CMs, but the risk for a BCVA loss of 3 Snellen lines or more or deterioration of BCVA to less than 20/200 (decimal scale, 0.1) is approximately 50% at 10 years.³³⁴ With proton beam radiation therapy, a nearly 100% local tumour control and satisfying long-term visual outcomes for tumours located further than 3 mm from the fovea and optic disc have been achieved after treatment of small tumours.^{335,336} The probability of patients retaining BCVA of 20/200 or better was approximately 50% at 10 years.³³⁵

6.4.3. Transpupillary thermotherapy

TTT was introduced in the Netherlands as a treatment modality for CMs in 1995.^{48,337,338} It was initially described in combination with Ru¹⁰⁶ brachytherapy in an attempt to avoid radiation-related complications and preserve BCVA by reducing the radiation dose.^{48-50,339} However, the reduction in loss of BCVA proved to be insignificant compared to treatment with Ru¹⁰⁶ alone.^{314,340} TTT is now additionally used as a primary therapy or adjuvant therapy after brachytherapy.³⁰⁷

In TTT, a diode laser beam is directed through the dilated pupil to the apex of the tumour for approximately one minute, resulting in an increase in the tumour temperature to 45-60°C. The consequence of this heating is a tumour necrosis to a maximum depth of 4 mm,^{337,338} which is therefore considered to be the size limit for the indication of TTT as a primary treatment.^{48,57,338,341-343} Tumour regression continues for months and leaves an atrophic chorioretinal scar. The importance of aiming for a completely flat tumour scar is to reduce the risk of a local recurrence, although recurrent tumours have been reported to arise from completely flat scars.^{55,56,333,344}

When a tumour is small and thin, TTT may be considered as a primary treatment provided that the patient accepts that such treatment may need to be followed by radiotherapy in cases of recurrent or persistent tumours.⁵³ TTT causes an immediate local scotoma in the visual field, and delayed effects include macular scarring, epiretinal membranes, retinal vascular occlusions, and vitreous haemorrhages.^{292,342,344}

High local recurrence rates, up to 45%⁵²⁻⁶⁰ have been a concern following TTT. Several studies have reported higher recurrence rates for parapapillary tumours compared to non-parapapillary lesions.^{57,59,337,342,345} The overlying retina undergoes atrophy, but the underlying sclera is resistant to hyperthermia.³³⁸ Consequently, TTT may be associated with intra- and extra-scleral recurrences, which are rare after brachytherapy.^{52,53}

6.4.4. Proton beam therapy

One option for radiotherapy is external beam radiation using charged particles such as protons²⁸³ despite facilities for proton beam only being available in a limited number of centres around the world. This technique allows for a uniform dose distribution to a circumscribed target volume including the tumour, resulting in improved local tumour control and less damage to healthy tissues.³⁴⁶ Proton beam therapy is generally preferred for larger tumours that are not eligible for brachytherapy or for posteriorly located tumours, in an attempt to spare the macula and optic disc.³⁴⁷

Local tumour recurrence rates for CM of patients with tumours of all sizes have been reported at approximately 4% at 5 years³⁴⁷ and 5% at 10 years.³⁴⁸ The local recurrence rate was 5% at 15 years for tumours with a median of 5.3 mm in thickness and 13.2 mm in LBD.³⁴⁹ For T1 tumours, no recurrences were observed during a mean follow-up of 10 years.³³⁵

BCVA outcomes following proton beam therapy depend on tumour thickness and proximity to the fovea and optic disc.^{335,350} Patients with a tumour located at least 3 mm from these posterior structures generally do not develop clinically significant radiation vasculopathy.³⁵¹ The risk for visual loss at 5 years has been estimated at 68% for small and medium-sized tumours.³⁵² Five-year cumulative rates of maculopathy and papillopathy were 64% and 35 % when the analysis was restricted to a group of patients with small- to moderate-sized tumours located within 4 disc diameters of the optic nerve or macula.³⁵² The frequency of losing BCVA of 20/200 for tumours located further than 3 mm from fovea and optic disc was approximately 7% at 5 years, and 18% at 10 years.³³⁵ The corresponding rates for tumours located closer than 3 mm from the posterior structures was approximately 40% at 5 years and 53% at 10 years.

6.5.5. Experimental treatments

Photodynamic therapy (PDT) with verteporfin was originally used for choroidal neovascularization in age-related macular degeneration.³⁵³ It has been reported that PDT induces tumour regression for 62% to 100% of treated small posterior CMs,^{354,355} resulting in treatment outcomes that are less reliable with regard to tumour regression compared to other treatment modalities³⁵⁶ but without the associated decrease in BCVA.^{355,357} Although PDT has been reported to be effective in both amelanotic and pigmented tumours,³⁵⁸ doubts have been raised concerning its efficacy in treating the pigmented tumours.^{359,360} PDT was additionally reported to be more effective for treating tumours with three or fewer risk features for growth.³⁵⁴

AU-011 is a novel virus-like particle-drug conjugate that selectively binds to cancer cells and enables a targeted therapy for UM.^{361,362} The drug is administered through intravitreal injection and subsequently activated with an infrared laser, which leads to targeted tumour cell necrosis while simultaneously sparing healthy tissue.³⁶³ Preliminary results support the further development of this modality.³⁶⁴

6.6. METASTASIS AND SURVIVAL

Metastatic UM is the leading cause of death in UM patients, with the mortality rate being higher than 50% after 25 years of primary treatment for medium-sized and large tumours.^{20,92,365,366} Small tumours have a much better prognosis, with a metastatic rate of approximately 12% in 10 years.²⁴ It is not known exactly when dissemination takes place. Clinically evident metastases are detected in less than 1–3 % of all patients at the time of UM diagnosis, and the median length of time until detection of metastasis is 2 years after treatment for UM in general.^{92,367} It has been estimated that tumours may have the capacity to disseminate years before the diagnosis of the primary tumour.^{190,368} Studies of tumour doubling times of metastatic lesions have suggested that metastasis may commence when the primary tumour continues to be small, that is approximately 3 mm in LBD and 1.5 mm in thickness.¹⁹³ The smallest CMs that have been reported to metastasize have been at least 1.7 mm thick³⁶⁹ and 5.0 mm in LBD^{156,292,369-375}. Growth of metastases may be delayed, and metastases may appear clinically up to three decades after successful treatment of the primary tumour due to a slow growth pattern and later cytogenetic progression.^{20,92,190}

UM has a predilection to metastasize to the liver in over 90% of cases.^{19,376,377} Liver is the only site in 33–56% of metastatic patients.^{5,92,366,376,378} Other less common, and generally secondary, sites are the lungs, subcutaneous tissue, bones, brain, adrenal glands, and the heart.^{19,376,379-381} Metastases are more widespread than clinically suspected in patients who undergo autopsy.³⁶⁶ Symptoms vary, depending on the affected organ; yet, more than 60% of patients are asymptomatic when their metastases are detected.³⁸²

When a six-monthly screening protocol is applied, metastases are detected before the onset of symptoms in more than 90% of UM patients.³⁸³ Different protocols have been reported, but a majority of UM patients are checked for liver metastases every 6 or 12 months, depending on their tumour stage, using imaging techniques such as abdominal US, MRI, and liver function tests (LFTs).³⁸²⁻³⁸⁵ New abnormalities in the LFTs or appearance of the liver are highly suggestive of metastasis. However, LFTs can be normal in approximately one-third of cases with liver metastasis.³⁸² Abdominal US combined with LFTs is a sensitive screening modality, particularly when followed by a confirmatory MRI in cases of newly detected lesions.³⁸⁶ A histopathological confirmation of suspected metastases is performed if possible – and certainly when active treatment is to be proposed – depending on the general condition of the patient.

In the presence of metastases, survival prognosis is poor as no curative treatment is generally available at the moment. The median survival of patients with metastatic UM ranges from 6 to 13 months.^{23,367,387,388} The death rate following clinically diagnosed dissemination is 80% at 1 year and 92% at 2 years, and a long-term survival time of more than 5 years is exceptional.^{19,389-391} Systemic treatment can only slow metastatic growth and improve survival to a limited extent.^{5,388}

Although there are many therapeutic options, there is no established standard of care for patients with metastatic disease. These options include systemic and intrahepatic chemotherapy, transarterial chemoembolization, isolated hepatic perfusion, protein kinase inhibitors, selective internal radiation therapy, immunoembolization, immunosuppressants, partial hepatectomy, microwave ablation, selective internal radiotherapy, and liver-directed thermotherapy.^{23,389} With a local resection of solitary liver metastases, a survival of seven years has been reported, but this is only available for a minimal number of selected patients.³⁹²⁻³⁹⁵ Patients with a loss-of-function mutation in the MBD4 gene are likely to benefit from CPI immunotherapy, a group of agents successfully used in cutaneous melanoma therapy.^{118,130-132}

7. Aims of the study

The first two studies (Study I and II) in this thesis analysed the treatment results of small CMs of less than 10 mm in LBD with the 10-mm Ru¹⁰⁶ plaque. The second study compared these results with those of similarly sized melanomas treated with the 15-mm Ru¹⁰⁶ plaques. The third study (Study III) reported sizes and clinical features of the smallest CMs that metastasized. The fourth study (Study IV) reported the growth rates and doubling times of presumed incipient CMs.

The specific aims of this study were as follows:

- I. To assess local tumour control, side-effects, and BCVA in a population-based consecutive series of patients who received brachytherapy for CM less than 10 mm in LBD with the 10-mm Ru¹⁰⁶ plaque.
- II. To assess local tumour control, side-effects, and BCVA in a population-based consecutive series of patients who received brachytherapy for CM less than 10 mm in LBD with the 10-mm or 15-mm Ru¹⁰⁶ plaque and compare the outcomes between two groups.
- III. To empirically determine the size limit at which a small CM acquires the capacity to metastasize and to assess the clinical characteristics of small fatal CMs
- IV. To analyse tumour doubling times and growth rates as potential alternative diagnostic criteria for incipient small CMs in a population-based study with a consecutive series of patients with presumed incipient CM that was subsequently managed with primary TTT

8. Material and methods

8.1. PATIENTS AND STUDY DESIGN

8.1.1. Studies I and II

Patients who were clinically diagnosed with a CM less than 10 mm in LBD and managed with primary brachytherapy with the 10-mm (Study I and II) or a 15-mm (Study II) Ru¹⁰⁶ plaque were eligible to participate in these studies. A patient was ineligible if the tumour involved the ciliary body or iris. The database of the Ocular Oncology Service, Department of Ophthalmology, Helsinki University Hospital, a national referral centre for UM patients, was reviewed.

The time period for Study I began in October 1998 – at which point the 10-mm plaque became a treatment of choice in Helsinki University Hospital – and ended in December 2010, with follow-up data being collected until the end of 2011. Within this time period, a total of 118 patients with an UM less than 10 mm in LBD were identified, of whom, 48 were treated with the 10-mm plaque. Three patients had a ciliary body or iris tumour, resulting in 45 patients being enrolled.

The time period of Study II began in October 1998 as well. Patients were treated with the 10-mm or 15-mm plaque until December 2014, and follow-up data were collected until the end of 2017. Within this period, a total of 234 patients with CMs less than 10 mm in LBD were identified, of whom, 76 were scheduled for treatment with the 10-mm and 88 were scheduled with the 15-mm plaque.

8.1.2. Study III

Data on consecutive patients diagnosed with a CM of 3 mm or less in thickness and 9 mm or less in LBD who subsequently developed metastases were collected from 10 ophthalmic oncology centres in Europe that agreed to participate in the study, including 9 patients from the Ocular Oncology Service, Department of Ophthalmology, Helsinki University Hospital. The data of 56 patients were collected, of whom, 11 were excluded after an eligibility check and 45 enrolled. Excluded were 5 patients who had tumours larger than the eligibility criteria, 3 patients who had incomplete key data, one patient who was diagnosed with pulmonary metastases from an epithelioid cell melanoma, and 2 patients who had only extrahepatic metastases without histopathological verification. The patients had been diagnosed with CM between 1962 and 2010.

8.1.3. Study IV

All 9 consecutive patients who had been diagnosed with a presumed incipient CM following documented growth and subsequently treated with primary TTT between 2010 and 2017 in

the Ocular Oncology Service, Department of Ophthalmology, Helsinki University Hospital were eligible for this retrospective study. One of the patients had a previous T2aN0M0, Stage IIA, CM in the same eye that had been treated with the 15-mm ruthenium plaque 2.5 years before the second tumour (enrolled in the present study) was diagnosed.

8.2. PRIMARY TREATMENT

8.2.1. Brachytherapy (Studies I and II)

The treatments were performed by three experienced ocular oncologists over the 16-year period. The prescription point was tumour thickness plus 1 mm for the sclera. Treatment times and doses were calculated from the central axis depth dose curve. The standard prescription dose was either 100 Gy or 120 Gy to the apex depending on the thickness of the tumour and aimed to deliver a minimum dose of 250 Gy to the sclera. Overlying extraocular muscles were disinserted if needed to ensure accurate positioning of the plaque in relation to the tumour.

The cord diameter of the radiation window of the plaque in spherical form is 9.4 mm and 12.9 for 10-mm and 15-mm plaques, respectively (Eckert & Ziegler BEBIG GmbH, Berlin, Germany). The radiation window is surrounded by a 0.75-mm-wide inactive border. The radiation window is surrounded by a penumbra of scatter radiation, which is used as justification for eccentric plaque placement in order to limit irradiation to adjacent tissues.^{310,396} Plaque position and distance to the tumour apex were verified on the first postoperative day with US in view of a calculation update of the treatment time. However, this practice was not in use in the early study years.

8.2.2. Transpupillary thermotherapy (Study IV)

TTT was delivered through a dilated pupil in periocular anaesthesia with an infrared diode laser (MedArt 426, MedArt, Hvidovre, Denmark) at a wavelength of 810 nm which was adapted to a slit lamp biomicroscope and which used beam diameters of 0.8 to 3.0 mm and a contact lens (Area Centralis or Quadraspheric, Volk Optical, Mentor, OH). The spot was applied 1 to 4 times with increasing power if needed (median, 1,500 mW; range, 700–3500) for a median duration of 120 s (60–285) to achieve a greyish white colour on the surface of the tumour. Overlapping spots, if needed, were used to cover the entire tumour, and there was a margin of 0.5 to 1.0 mm on all sides. TTT was repeated 1 to 5 times at intervals of 2–7 months with a goal of achieving a white chorioretinal scar. Three ocular oncologists performed all treatments.

8.3. DATA COLLECTION

8.3.1. Clinical evaluation

8.3.1.1. Studies I and II

The tumour size was primarily determined by indirect ophthalmoscopy and B-scan US. Biomicroscopy and indirect ophthalmoscopy were used to localize the tumour margins in order to measure the distance from the posterior tumour margin to the optic disc and foveola and evaluate the presence of SRF and OP. Patients were treated if they had a presumed small CM that either had been observed to grow or had high-risk features for growth^{27,42,397} (Tables 3–4). The BCVA was measured using a test-type projector (Rodavist 2 and 524; Rodenstock GmbH, Ottobrunn, Germany) in even decimal steps from 20/20 (decimal scale, 1.0) to 20/200 (decimal scale, 0.1) plus 20/400 (decimal scale, 0.05). A BCVA worse than 20/400 was recorded as counting fingers at 2 m, 1 m, and 0.5 m; hand movement; light perception; and no light perception.

Table 3. Risk factors for growth^{27,42} in 45 (Study I) and 164 (Study II) patients with small choroidal melanomas less than 10 mm in largest basal diameter, of whom 76 were managed with the 10-mm plaque and 88 with the 15-mm plaque (Studies I and II). Study I patients are included in Study II.

Risk Factors	Study I				Observed Growth, n (%)			
	Study I		Study II, n (%)		Study I		Study II, n (%)	
	n (%)		10-mm	15-mm	n (%)		10-mm	15-mm
	n=45	n=76	n=88	P*	n=18	n=29	n=32	P*
Thickness	18 (40)	30 (39)	63 (72)	<0.001	6 (33)	6 (21)	19 (59)	0.004
>2 mm								
Subretinal fluid	28 (62)	48 (63)	55 (63)	0.99	5 (28)	8 (28)	11 (34)	0.81
Symptoms	25 (56)	40 (53)	39 (44)	0.35	5 (28)	8 (28)	5 (16)	0.35
Orange pigment	28 (62)	47 (62)	45 (51)	0.21	7 (39)	13 (45)	11 (34)	0.44
Margin								
touching disc	3 (7)	3 (4)	1 (1)	0.34	1 (6)	1 (1)	0 (0)	0.48
within 2 DD	26 (58)	40 (53)	41 (47)	0.53	5 (28)	8 (28)	14 (44)	0.19

DD = disc diameter, *Fisher's exact test

Table 4. Number of risk factors⁴² for growth in 45 (Study I) and 164 (Study II) patients with small choroidal melanoma less than 10 mm in largest basal diameter who were treated with the 10-mm and the 15-mm ruthenium¹⁰⁶ plaque. Study I patients are included in Study II.

Number of Risk Factors	Study I n (%)	Study II, n (%)		P*
	10-mm n=45	10-mm n=76	15-mm n=88	
0	4 (9)	9 (12)	5 (6)	0.12
1	5 (11)	9 (12)	10 (11)	
2	9 (20)	16 (21)	17 (19)	
3	11 (24)	14 (18)	30 (34)	
4	11 (24)	19 (25)	21 (24)	
5	5 (11)	9 (12)	5 (6)	

*non-parametric test for trend

In Study I, 4 (9%) patients who had none of the risk features were treated due to observed growth. In Study II, 14 patients had a tumour without risk features, of whom, 12 (7%) were treated because of observed growth, one had a tumour that had not been seen during regular glaucoma follow-up, and one had a ruptured Bruch's membrane.

8.3.1.2. Study III

This retrospective study was conducted using patient charts, archival images, and pathology data. Members of the European Ophthalmic Oncology Group submitted anonymous data from 10 ocular oncology services to a secure survey website. The data requested included date of birth, diagnosis, primary treatment, local recurrence, and diagnosis of metastases as well as sex, ethnicity, involved eye, BCVA, history of a previous naevus, high-risk features for growth and metastasis,^{27,42} observation before treatment, type of the primary treatment, histopathologic diagnosis, secondary treatments, systemic treatment, and last status.

8.3.1.3. Study IV

Digital fundus images were used to localize the tumour margins in order to estimate the distance from the posterior tumour margin to the optic disc and foveola and to evaluate its growth as well as the presence of SRF and OP. Based on OCT, a tumour was considered flat if the tumour area was not elevated compared to the adjacent normal choroid. The BCVA was measured using a test-type projector (Rodavist 2 and 524; Rodenstock GmbH, Ottobrunn, Germany) in even decimals from 1.0 (20/20) to 0.1 (20/200).

Diagnosis was based on verified tumour growth relative to the age of the patient and reported chance of having a growing choroidal nevus, TDT, growth rate, and to the calculated age of tumour origin. Patients were selected to have primary TTT instead of brachytherapy due to the small tumour sizes.

The initial data were collected from the medical records of each patient, including details on age, race, gender, diabetes, ocular history, tumour growth, high-risk factors, BCVA, tumour location, tumour measurements, and information from TTT. Data after completion of TTT included final BCVA, tumour control, TTT complications, and last status.

8.3.2. Calculations of growth rates and doubling times

For this study, the tumour growth rate was calculated starting from its LBD and using a theoretical measure of 0.05 mm when the tumour was not visible in the first photograph. The tumour volume was then estimated presuming a cylindrical form because all tumours were assumed to have the same thickness as the choroid since the tumours did not thicken the adjacent choroid on the OCT images. The surface area of the tumour was calculated by using ImageJ software (National Institute of Health, Maryland, USA, Version 1.51). Two investigators measured the tumour area independently by two different methods (manual drawing and automatic recognition of tumour margins) and the final area that was taken was the mean of these four measurements. The date of the diagnosis of the primary tumour was the day on which an ophthalmologist first made the diagnosis of CM due to verified growth. The date of first observation was the date when the lesion had been first documented. Observation time (T) was the difference between these two dates.

TDT was calculated according to Schwartz³⁹⁸:

$$\text{TDT} = \frac{T}{\left(\frac{\log_{10} \left(\frac{V_{dg}}{V_{ini}} \right)}{\log_{10}(2)} \right)} \quad (\text{equation 1})$$

where T = observation time, Vdg = tumour volume at diagnosis, and Vini = initial tumour volume.

The predicted initial age was additionally calculated to estimate whether the tumours had originated in adulthood and thus, were likely to be *de novo* tumours (equation 2). A theoretical initial LBD of 0.05 mm was used when a tumour likely would have been visible.

$$\text{Age}_{\text{pred}} = \text{Age}_{\text{ini}} - \frac{\text{TDT} \times \frac{\log_{10} \left(\frac{V_{ini}}{V_{orig}} \right)}{\log_{10}(2)}}{365.25} \quad (\text{equation 2})$$

where Age_{pred} = age when the tumour was predicted to be visible, Age_{ini} = age when the tumour was initially observed, TDT = tumour doubling time (equation 1), Vini = initial tumour volume, and Vorig = theoretical volume 0.000125 mm³ based on 0.05 mm diameter. Calculations were based on a presumed constant exponential growth.³⁹⁹

For comparison, TDTs collected for 196 patients with a UM reported in 4 articles to serve as reference data.^{191,192,400,401} Omitting two outliers with exceptionally long TDT,¹⁹¹ the median

UM TDT was 521 days (range, 10–8766). The reported median growth rate for choroidal naevi, that is 0.04 mm/year (range, 0.01–0.2) and 1.1 %/y (range, 0.2–11.6), for 89 naevi that showed photographic evidence of enlargement⁴³ – excluding a tumour that later had been diagnosed as a CM – was used as a counter reference.

8.3.3. Assessment of outcomes (Studies I and II)

Patients were followed at 3, 6, and 12 months; then every 6 months until 3 to 4 years after brachytherapy, and annually thereafter. Biomicroscopy, indirect ophthalmoscopy, fundus photography, and B-scan echography (I3 System-ABD; Innovative Imaging, Sacramento, California) were used to assess tumour control. Recurrences were coded as being vertical, marginal, diffuse, or extrascleral.⁴⁰²

The location of the plaque was retrospectively modelled in relation to the tumour to facilitate determining the reasons for a local recurrence. The location of the plaque was coded as ‘covering the entire tumour with a safety margin’ (covered), as ‘being parallel’ (eccentric), or as ‘not covering the entire tumour’ (not covered). A circle was digitally drawn to the post-treatment photographs, that corresponded in size to the radiation window. This was done using radiation and diathermy scars, with the presumption that the diameter of the optic disc was 1.5 mm.

Complications were assessed, as outlined, in a previous study on brachytherapy from the same ocular oncology service.⁴⁰³ Radiation maculopathy was diagnosed based on the appearance of retinal haemorrhage, microaneurysms, microinfarcts, oedema, exudation, and a decrease in BCVA. A tumour recurrence involving the fovea, scarring, or a macular pucker were categorized as ‘maculopathy for other reasons’. Optic neuropathy was categorized as radiation optic neuropathy based on the appearance of oedema, peripapillary splinter haemorrhages, or microinfarcts.

8.3.4. Assessment of survival and metastatic status (Studies I and II)

LFTs and upper abdominal US were performed at the time of diagnosis and annually thereafter, and a chest radiogram was performed only at diagnosis. Conspicuous findings of concern were verified with CT or MRI and, eventually, histopathologically. Causes of deaths were verified from death certificates obtained with permission from Statistics Finland, autopsy reports, and patient charts.

8.4. STATISTICAL METHODS AND DATA ANALYSIS

Follow-up data were collected prospectively into a dedicated database (MS Access; Microsoft, Seattle, WA) and analysed with Stata statistical software (Release 10, 13 and 15; Stata Co, College Station, TX). A cumulative incidence analysis⁴⁰⁴ was used that accounted for competing events in order to analyse the time to individual complications and death

– except in the third study where the analysis of time to systemic metastasis and survival was based on the Kaplan-Meier product-limit method since, by definition, all patients developed systemic metastases and none died of other causes.

All tests were 2-tailed, and a P-value less than 0.05 was considered statistically significant. The data were summarized as frequencies and percentages for categorical variables and as a median with a range for continuous variables. A Fisher's exact test was used to compare dichotomous variables. A non-parametric test for trend was used exploring the differences between two independent groups for ordinal variables. A Kruskal-Wallis test was used between two or more independent groups for continuous variables, and a Gray's k-sample test⁴⁰⁵ was used to compare cumulative incidences between two groups. Parameters including the time to local recurrence, low vision (BCVA < 20/65; decimal scale, 0.3), and blindness (BCVA < 20/400; decimal scale, 0.05) were modelled by univariable and multivariable Cox proportional hazards regression.

9. Results

9.1. TREATMENT WITH THE 10-MM OR 15-MM PLAQUE (STUDIES I AND II)

9.1.1. Baseline patient, eye, and tumour characteristics

All tumours treated with the 10-mm plaque were classified as T1aN0M0, stage I, by the AJCC TNM-System. Eighty-two percent of tumours treated with the 15-mm plaque were classified as T1aN0M0, stage I, and the remaining tumours with a thickness over 3 mm and LBD over 9 mm were classified as T2aN0M0, stage IIA (Fig. 3).¹⁴⁵

The melanomas of the patients treated with the 15-mm plaque were thicker as a group (median, 1.9 vs 2.6 mm, $P<0.001$), had larger LBDs (median, 7.1 vs 8.6, $P<0.001$) and longer distances from the posterior tumour margin to the center of the fovea (median, 2.0 vs 2.8, $P<0.001$) (Table 5, Fig. 3). In Study II, the median age at diagnosis (65 vs. 63 years; $P=0.59$), the median follow-up time for patients who were still alive (8.1 years, range 3.0–18.8 vs. 8.9 years, range 1.1–17.9; $P=0.67$), and the median BCVA (20/25 vs. 20/30; $P=0.54$) were comparable between the two groups (Table 5).

Table 5. Baseline characteristics at the time of diagnosis of 45 (Study I) and 164 (Study II) patients with a small choroidal melanoma less than 10 mm in largest basal diameter irradiated with the 10-mm (Studies I and II) or the 15-mm (Study II) ruthenium¹⁰⁶ plaque. Study I patients are included in Study II.

Characteristic	Study I median (range)	Study II, median (range)		P^*
	10-mm n=45	10-mm n=76	15-mm n=88	
Age, years	61 (26–88)	65 (26–88)	63 (27–88)	0.59
Follow-up time, years	5.2 (0.3–11.9)	8.1 (3.0–18.8)	8.9 (1.1–17.9)	0.45
Thickness, mm	1.9 (0.4–5.2)	1.9 (0.4–5.2)	2.6 (0.3–5.7)	<0.001
LBD, mm 7.0 (3.3–9.6)	7.1 (3.3–9.6)	8.6 (4.1–9.9)	<0.001	
Distance to foveola, mm	2.0 (0–8.5)	2.0 (0–13.0)	2.8 (0–13.5)	<0.001
Distance to optic disc, mm	3.0 (0–7.5)	3.0 (0–13.5)	3.0 (0–13.5)	0.79
BCVA	20/25	20/25	20/30	0.54

BCVA = Best corrected visual acuity, LBD = Largest basal diameter, * Kruskal Wallis test




Figure 3. 45 (Study I, red symbols) and 164 (Study II) small choroidal melanomas less than 10 mm in largest basal diameter (LBD) irradiated with the 10-mm (circle, red and black) or the 15-mm (square black) ruthenium¹⁰⁶ plaques. A, Scatterplot of radiation dose to the sclera against dose to the tumour apex. Dashed lines indicate minimum prescription doses. B, Scatterplot of LBD against tumour height. Long and short dashed lines enclose tumours which fulfil the American Joint Committee for Cancer (AJCC) Tumour, Node, Metastasis (TNM, 8th Edition)¹⁴⁵ and the Collaborative Ocular Melanoma Study (COMS)³¹ criteria for a small UM, respectively. C, Scatterplot of proximity of tumour margin to the foveola against proximity to the optic disc. Dashed lines indicate a distance of 2 DD (3 mm) from these structures, frequently considered to be a safety margin for not developing radiation maculopathy and optic neuropathy from ruthenium¹⁰⁶ brachytherapy. A–C, Solid and open symbols indicate tumours with and without recurrence, respectively.

9.1.2. Plaque positioning

The location of the plaque was assessed according to the surgical charts and by retrospective modelling (Table 6). In the comparative Study II, tumour coverage was statistically significantly different between the two groups ($P=0.002$) but differed to a lesser extent ($P=0.052$) after remodeling.

Table 6. Location of the plaque according to the surgical charts and by retrospective modelling of 45 (Study I) and 164 (Study II) patients with a small choroidal melanoma less than 10 mm in largest basal diameter irradiated with the 10-mm (Studies I and II) or the 15-mm (Study II) ruthenium¹⁰⁶ plaque.

Location	Surgical chart, n (%)				Modelling, n (%)			
	-----				-----			
	Study I	Study II			Study I	Study II		
	10-mm	10-mm	15-mm	P*	10-mm	10-mm	15-mm	P*
Covered	13 (29)	25 (33)	50 (57)	0.002	6 (13)	33 (43)	35 (40)	0.052
Eccentric	20 (44)	28 (37)	28 (32)		13 (29)	13 (17)	29 (33)	
Not covered	10(22)	22 (29)	10 (11)		18 (40)	24 (32)	16 (18)	
Indeterminate	2 (4)	1 (1)	0		7 (16)	6 (8)	7 (8)	

*non-parametric test for trend

9.1.3. Local tumour recurrence and survival

A total of 4 patients developed a local recurrence in Study I. A local recurrence was detected in 9 and 16 patients treated with the 10-mm and the 15-mm plaque, respectively, in Study II. Recurrences were controlled with a secondary iodine plaque, except in three patients, of whom one had been retreated with a secondary Ru¹⁰⁶ plaque and two had been treated with TTT. In Study II, the cumulative incidence of developing a local recurrence was comparable between the two groups (Table 7). By univariable Cox regression, a local recurrence was associated with an eccentric plaque location (HR 3.40 for a plaque not covering the tumour with a safety margin, $P=0.024$) (Table 6, Study II).

Of the 45 patients in Study I, 6 died during the follow-up. Five patients died for reasons other than metastatic disease, and one death was classified as being the result of possible metastasis to the brain as the only site, although this had not been histopathologically verified. Within the Study II period, a further 3 patients died of metastatic CM after a local tumour recurrence following treatment with the 10-mm plaque. Seven patients, of whom 4 had experienced a local tumour recurrence, died of metastatic CM after treatment with the 15-mm plaque. The development of metastases, all-cause, and melanoma-related mortality rates were comparable (Table 7). The time from treatment to metastases (median, 4.6 vs 4.5

years; $P=0.85$) and survival after diagnosis of metastases (median, 3.6 vs 6.1 years; $P=0.73$) were comparable between the two groups.

Table 7. Summary of outcomes using cumulative incidences after treatment of 164 patients with small choroidal melanomas less than 10 mm in largest basal diameter with the 10-mm or the 15-mm ruthenium¹⁰⁶ plaque

Outcome	Percentage at 5 years, (95% CI)			Percentage at 10 years, (95% CI)		
	Study II					
	Study I 10-mm	10-mm	15-mm	10-mm	15-mm	P^*
Recurrence	9 (3–20)	9 (4–17)	13 (7–20)	13 (6–21)	15 (8–23)	0.31
Metastases		4 (1–10)	6 (2–13)	4 (1–10)	10 (4–19)	0.50
All-cause mortality	11 (4–26)	10 (4–18)	9 (4–17)	24 (13–37)	29 (19–40)	0.24
Melanoma-mortality		3 (1–8)	2 (0.5–7)	3 (1–8)	8 (3–15)	0.33
Maculopathy						
Radiation	28 (15–42)	19 (11–28)	18 (11–27)	24 (15–35)	22 (14–32)	0.99
Other	15 (6–27)	4 (1–11)	1 (0.1–6)	10 (4–21)	10 (4–19)	0.54
Optic neuropathy	3 (0–12)	1 (0.1–7)	8 (3–15)	1 (0.1–7)	8 (3–15)	0.054
Low vision	17 (7–31)	14 (7–24)	24 (15–34)	21 (11–33)	30 (20–41)	0.36
Blindness	3 (2–12)	3 (0.6–10)	6 (2–13)	14 (6–25)	13 (6–23)	0.90

CI = Confidence interval, *Gray's k-sample test

9.1.4. Radiation maculopathy and optic neuropathy

Radiation maculopathy and optic neuropathy were the most frequent radiation-related complications. Radiation maculopathy was diagnosed in 14 eyes within the Study I period and in 17 and 20 eyes after treatment with the 10-mm and the 15-mm plaque in Study II, respectively. The cumulative incidence of developing radiation maculopathy and maculopathy other than radiation maculopathy were comparable between the two groups in Study II (Table 7). Additionally, the distance of the tumour margin to the center of the foveola (median, 0.5 vs 1.0 mm, $P=0.23$) was comparable in eyes that developed radiation maculopathy.

One eye developed radiation optic neuropathy after treatment with the 10-mm plaque (Studies I and II) of a tumour located at 4 mm from the optic disc. Optic radiation neuropathy developed in 7 eyes after treatment with the 15-mm plaque. The cumulative incidence rates of developing optic neuropathy (Table 7) and the distance of the posterior tumour margin to the margin of the optic disc (median, 4.0 vs 2.0mm, $P=0.12$) were comparable.

9.1.5. Preservation of visual acuity

A tumour thickness of more than 3.0 mm was associated with more frequent loss of vision, particularly after treatment with the 10-mm plaque, although the number of patients with such a thick tumour was small. Instead, a location of the tumour margin at less than 1.5 mm from the optic disc and foveola was a more prominent indicator for vision loss after treatment with the 15-mm plaque (Fig. 5, Study II). The cumulative incidences of losing a minimum BCVA of 20/65 (low vision) and the development of blindness (BCVA < 20/400) were comparable between the two groups (Table 7).

By univariable Cox regression, low vision and blindness of the tumour eye were associated with a shorter distance to the foveola, a worse baseline BCVA, and larger tumour thickness. Of bivariable models using plaque size, the model including the distance to the foveola independently and best predicted low vision (HR, 0.64 for each 1 mm increase in distance, $P < 0.001$; Table 8, Study II). Distance to the foveola was further associated with blindness in a similar bivariable model (HR, 0.68 for each 1 mm decrease in distance, $P = 0.004$).

9.2. SMALL METASTASIZING CHOROIDAL MELANOMAS (STUDY III)

9.2.1. Tumour Size and Characteristics

The median thickness of a SFCM at the time of treatment was 2.4 mm (range, 1.0–3.0 mm, Table 8) and its median LBD was 7.3 mm (range, 3.0–9.0 mm). None of the 45 tumours was less than 3.0 mm in LBD at the time of treatment (Fig. 1, Study III), whereas 27% of the 45 tumours ranged from 3.0–6.0 mm and 73% from 6.1–9.0 mm in LBD. Risk factor frequencies (Table 8) at the time of diagnosis were similar to those reported in a study on 35 small melanocytic choroidal tumours that metastasized²⁷ (Table, Study III).

Table 8. Baseline characteristics and outcomes of 45 patients with a small choroidal melanoma less than 3 mm in thickness, less than 9 mm in largest basal diameter, and who developed metastatic disease.

Age (years)*	57 (26–81)
Treated immediately	31 (69)
Observed for growth	14 (31)
BCVA*	20/40 (counting fingers–25/20)
Colour	
Dark brown	16 (36)
Light brown/amelanotic	24 (53)
Mixed	3 (7)
Distance (mm) (26/45)	
to the foveola*	0.8 (0–5.0) (26/45) †
to the optic disc*	1.0 (0–4.3) (26/45) †
Tumour location	
Temporal	24 (57)
Nasal	18 (43)
Tumour thickness (mm)	2.4 (1.0–3.0)
LBD (mm)	7.3 (3.0–9.0)
US hollowness	26/3/2‡ (31/45) †
Risk features	
over 2 mm in thickness	27 (60) (27/45) †
subretinal fluid	20 (63)
symptoms	37 (84)
orange pigment	21 (57)
margin within 2 DD	24 (92)
margin touching	7 (27)
Primary treatment	
Enucleation	15 (33)
Brachytherapy	18 (40)
Ruthenium ¹⁰⁶	13
Iodine ¹²⁵	5
Proton beam	11 (24)
TTT	1
Histopathology (20/45) †	
Epithelioid cells	15 (75)
Spindle cells	3 (25)
Extrascleral extension	1
Last status	
Died of metastases	37 (82)
Alive with metastases	6 (13)
Lost to follow-up	2 (4)

BCVA = Best corrected visual acuity; DD = Disc diameter; Data are no. (%) unless otherwise indicated; *Median (range); † (Number of events/numbers at risk); ‡ the acoustic profile lower/equal/higher than the surrounding choroid

9.2.2. Local recurrence, development of metastases, and survival

Local recurrence only occurred after conservative treatment. The cumulative incidence of developing a local recurrence was 17% (95% CI, 7–29) by 5 years and 19% (95% CI, 9–32) by 10 years. By study design, all patients presented metastatic disease. The cumulative incidence of developing metastasis was 51% (95% CI, 36–64) by 5 years and 85% (95% CI, 71–92) by 10 years after primary treatment. The Kaplan-Meier estimate of melanoma-related death following the detection of metastasis was 52% (95% CI, 37–66) at 1 year, 77% (95% CI, 60–88) at 2 years, and 84% (95% CI, 66–93) at 3 years.

9.3. INCIPIENT CHOROIDAL MELANOMAS (STUDY IV)

9.3.1. Baseline characteristics

The median age of the patients was 55 years (range, 41–77) when the tumour was first documented. All tumours were first detected in fundus photographs taken for various reasons (Supplemental text, Study IV). Two patients had fundus photos without a visible tumour. The median tumour thickness was 0.20 (range, 0.15–0.29) mm on OCT. None of the tumours was thicker than the adjacent choroid. The median LBD was 0.52 (range, 0.05–1.09) mm. None of the tumours showed any risk features for malignancy when first documented.

9.3.2. Tumour characteristics at diagnosis

The median time to observe tumour growth before its diagnosis as an incipient CM was 3.3 years (range, 2.2–7.3 years), at a median age of 57 years (range, 47–78) (Fig. 4A). Tumour thickness was unchanged, but the LBD had increased to a median of 1.6 (range, 0.93–2.32) after this observation. All patients had a BCVA of 20/25 (decimal scale, 0.8) or better in the tumour eye, except one patient with a cataract who had a BCVA of 20/40 (decimal scale, 0.5). The median distance to the foveola was 3.7 mm (range, 0.5–5.3) and 2.3 mm (range, 0–6.8) to the optic disc margin (Fig. 4B). The posterior tumour margin touched the optic disc in one eye and extended to within 2 DD of the disc margin in six eyes. One patient had developed four risk features: symptoms, SRF, OP, and tumour margin touching the optic disc. Eight tumours grew symmetrically, including the 2 in the eyes in which a previously normal fundus had been documented. Two patients had a presumed previous nevus that started to grow, of whom one had an asymmetrically growing tumour.

The median growth rate was 0.25 (range, 0.11–0.72) mm/y corresponding to 34 %/y, which was faster than the growth rate derived from the reference data of 88 growing naevi⁴³, ($P < 0.001$, Kruskal-Wallis test) they were compared with. Restricted to the 7 *de novo* tumours, the corresponding rates were 0.29 mm/y (range, 0.15–0.72) and 55 %/y (range, 24–1437).

The tumour volume doubled at a median of 2.4 (range, 0.95–9.6) times. The median TDT (Fig. 4A) of 609 days (range, 97–1612) for all tumours, and 393 days (range, 97–609) for *de novo* tumours was comparable with the reference data of 194 melanomas^{191,192,400,401} ($P=0.62$, Kruskal-Wallis test).

Excluding the two likely transformed naevi, the predicted interval from the time when the tumour would have been invisible and the corresponding predicted theoretical patient age was 11.7 years (range, 0.6–34.6) and 51 years (range, 32–63), respectively.




Figure 4. Nine small choroidal melanomas treated with primary transpupillary thermotherapy. **A**, Scatterplot of observation time against tumour doubling time. **B**, Scatterplot of proximity of tumour margin to the foveola against proximity to the optic disc. Solid and open symbols indicate previous naevi and *de novo* melanomas, respectively. Red symbols indicate the tumour of the patient who had a previous T2a, IIa choroidal melanoma.

9.3.3. Transpupillary thermotherapy

One to four (median, 1) TTT applications were applied. Based as well on OCT, the treatment outcome was a white scar with minor RPE remnants in 56% of the patients, a white scar with a focal reactive proliferation of the RPE in 33%, and, as a result of an RPE tear, a scar with prominent reactive RPE proliferations in 1 patient. The median follow-up time after the last TTT session was 2.1 years (range, 7.5 months–8.7 years). None of the patients had developed a local tumour recurrence or metastases within the study period. The BCVA was unchanged in all but two eyes. One patient with a tumour under the papillomacular bundle had a BCVA of 20/40 (decimal scale, 0.5) after completion of TTT. She developed a macular epiretinal membrane and underwent vitrectomy 8 years after TTT. The second patient had a BCVA of 20/125 (decimal scale, 0.16) even after cataract surgery, because of a non-exudative age-related macular degeneration.

10. Discussion

10.1. TREATMENT WITH THE 10-MM OR THE 15-MM PLAQUE

Tumours treated with the 10-mm plaque were more posteriorly located, and the plaque less frequently covered the entire tumour with a safety margin, making these tumours theoretically more vulnerable to recurrences. The observed recurrence rates were low^{307,314,315,320,406} and comparable after treatment with both plaques, despite the fact that the 10-mm plaque emits less scattered radiation. While tumours treated with the 15-mm plaque were larger, neither tumour size nor dose to apex was associated with local recurrence in multivariable modelling. An eccentric plaque location was a risk factor for a local tumour recurrence with both plaques, unlike the conclusion in the original paper that reported on the eccentric technique.³¹⁰

TTT has been associated with a relatively high risk of local recurrence with local recurrence rates between 7–45%.^{56-59,292} The Kaplan-Meier estimate for tumour recurrence after TTT has been 26% in 5 years⁵³ compared to a corresponding 9% rate after brachytherapy with the 10-mm plaque in Study I. The number of risk features for growth did not predict local recurrence after treatment with the 10-mm plaque, unlike after TTT.⁵⁶ Previous reports advise that small CMs with multiple risk features should be treated with methods other than TTT.⁵³

Tumour location likely contributed to tumour recurrences after treatment with the 10-mm plaque. Round eccentric plaques must be used with caution³¹⁰ when managing CMs which are located within 1 DD from the optic disc, where it is possible that a plaque might tilt along the optic nerve sheath. Three of four recurrent tumours in Study I were located less than 1.5 mm from the optic disc. Furthermore, peripapillary tumours have been found to be at a high risk of local recurrence also after TTT.^{56,292}

The frequent perifoveal location in these studies explains the high rate of developing radiation maculopathy and thus the effect on reading vision. The unadjusted rate of developing radiation maculopathy was equal with both plaques (19% and 18% at 5 years), as estimated by cumulative incidence analysis, and thus lower than in Study I (28%). The factors associated with a decrease in BCVA were tumour thickness over 3 mm, LBD over 7 mm, and tumour location less than 1.5 mm from the foveola. Although the loss of a minimum vision of 20/65 (=low vision) and 20/400 (=blindness) vision was comparable between eyes treated with the 10-mm and 15-mm plaques overall, multivariable analysis showed that the distance to the foveola, which differed between the two groups, was a factor affecting the risk of vision loss. Controlling for this distance, the risk of losing vision was higher when treating with the 15-mm plaque.

Melanoma-related mortality after treatment with the 10-mm or the 15-mm plaque was not higher than the estimate for small (T1a, stage I) tumours in general²⁴ despite the fact that a majority of tumours irradiated with the 15-mm plaque were T2a, stage IIA. The 2% melanoma-related mortality in Study I was comparable to that of TTT studies, where metastases developed in up to 5% of patients.^{56,57,292,332}

In the literature, a 100% local tumour control has been reported after brachytherapy with the gamma ray emitter Pd¹⁰³ for tumours less than 10-mm in LBD.⁴⁰⁷ However, the offset was a higher rate of radiation maculopathy in 43% of eyes and radiation optic neuropathy in 21% of eyes irradiated with Pd¹⁰³.

Comparable local tumour control and survival rates, as well as a low rate of complications other than radiation maculopathy and relatively good visual outcomes, might favour the treatment of small CMs less than 10 mm in LBD with the 10-mm Ru¹⁰⁶ plaque over the 15-mm plaque. The local recurrence rate after treatment with the 10-mm plaque appears to be lower compared with TTT^{53,56-59,292} and not worse than with the 15-mm plaque when managing patients with a CM less than 10-mm in LBD. Vision loss was more probable after the 15-mm plaque when the tumours were located close to the foveola.

10.2. SMALL FATAL CHOROIDAL MELANOMAS

A total of 3 patients with a small CM of less than 5 mm in LBD who developed metastases were found in the data collected from 10 ocular oncology services in Europe. A CM that metastasized and was less than 3 mm in LBD was not found, which suggests that 3.0 mm could be the limit from which a CM first gains its ability to metastasize, supported by the finding that no tumour smaller than 3 mm in diameter has been reported to metastasize, and the fact that the participating centres did not find any such tumour in a total of over 10,000 treated melanomas. This limit would, however, be subject to challenge and modification if necessary.

This multicentre study further suggests that if the diagnosis cannot be otherwise obtained, it is safe to observe choroidal tumours until they reach 3 mm in LBD in order to verify growth. This will push the size limit for early detection and treatment. Three of the tumours in this study grew over the 3 mm in LBD while being observed, and, on one hand, one can speculate whether metastasis in these patients could have been prevented by earlier treatment. On the other hand, all tumours, once reasonably diagnosed as a CM, regardless of their size, should not continue to be observed but should be treated.

Four percent of the patients in this study did not present high-risk features for growth, and the frequencies of these criteria at the time of their confirmed CM diagnosis were similar to those in earlier reports on small melanocytic choroidal tumours.²⁷ A comparison of the number and type of risk features for growth between the SFCMs in Study III with those

reported for the unselected CMs in the same size range in Study I suggested that there was no difference. Additionally, the number of risk features for growth has not been associated with a specific GEP class predictive of patient survival.⁴⁰⁸ Consequently, according to both this present study and the findings from the literature, no presently known clinical tumour characteristics can predict the development of metastases.

10.3. DIAGNOSIS OF INCIPIENT CHOROIDAL MELANOMAS

All tumours were flat and 2 mm or less in LBD at the time of treatment, which would most likely correspond to the tumour size of a naevus.³¹ Only one patient had developed a tumour with 4 high-risk features, and another patient had a tumour touching the disc margin as its only risk feature.^{27,42} The diagnosis was based on a verified faster tumour growth compared with that of growing naevi.⁴³ As these incipient tumours were too small for a biopsy,^{149,209-215} they could only be reasonably diagnosed by observed growth. The growth rates reported in this study were higher than those calculated from the reference data of naevi, while the TDTs were comparable with those from the reference data of melanomas, supporting the melanoma diagnosis of these tumours. The patient who had a history of a previously treated ipsilateral CM had the shortest TDT, and, therefore, the possibility of a local tumour metastasis could not be excluded.

It is generally presumed that choroidal naevi emerge during puberty, and that, consequently, new ones should not appear during adult life. For this study, the theoretical age at which each tumour diagnosed as an incipient CM would not have been visible was calculated. With one exception, this age result was 32 years or older. In the reference group from the literature, none of the growing naevi had developed new risk features.⁴³ Growth of these naevi had been observed in, respectively, 54%, 34%, and 19% of the patients younger than 40 years, 41 to 60 years, and older than 60 years. Five out of the nine patients in Study IV were 47 to 56 years old, while the other four belonged to the oldest group. One of the reference studies including 217 patients with a CMs less than 3.5 mm in thickness identified a patient age over 60 years as a new risk feature for predicting a GEP class 2.¹⁴⁹

A high local recurrence rate is a concern after primary TTT treatment of CMs when compared with radiation therapy.⁵² However, such a recurrence in these incipient tumours would likely be less than 3 mm in LBD and therefore be highly unlikely to metastasize.⁶¹ However, because recurrences have been reported to develop a decade after TTT,^{53,56} this study recommends a long-term regular follow-up for these patients, as suggested by other investigators.⁵³

A further recommendation can be made that all naevi should be documented, involving mandatory regular check-ups to observe possible changes in lesion sizes. Risk features for the malignancy of choroidal lesions have been reported; however, these features generally develop later in the evolution of the tumour.^{4,103} Currently, there is no effective treatment for

metastatic disease, and it is, therefore, necessary to find ways to diagnose and treat patients with CM as early as possible to prevent the occurrence of this fatal event.

10.4. STUDY STRENGTHS AND LIMITATIONS

The Helsinki University Hospital, Department of Ophthalmology, in which this study was conducted, is a national referral center for UMs that offers nationwide data. However, the late follow-up of patients coming from outside the Helsinki region is accomplished in other university hospitals or central hospitals starting 3 or 4 years after treatment if there is no evidence of recurrence. A limitation of all these retrospective studies is that a few patients had partially missing data for their individual visits to other university or central hospitals during their later follow-ups. Additionally, the number of patients in all the studies was small because of the rarity of this disease and the Finnish population amounting to only 5.5 million individuals. This small data set was compensated for by the fact that the present series were consecutive and population-based.

Study II was not randomised, as the plaques were not chosen systematically, and, therefore, the two groups were not entirely comparable. A multivariable analysis was used to adjust for these differences, which aimed at reducing bias. Small CMs are uncommon, making it difficult to organize a prospective study with an adequate sample size or, further, a randomised one.

Conservatively treated patients in all the studies neither received histopathological nor genetic verification, which, however, was in line with the usually reporting for these smallest tumours.

When comparing the results of this thesis for Study I and IV with the reference data in the literature, it is important to take into account the many differences both in inclusion criteria and definitions of complications. A longer follow-up is needed in Study IV to confirm the complete tumour control with TTT.

10.5. FUTURE ASPECTS

It would be ideal to aim for the earlier diagnosis of CM with a diagnostic accuracy of 100% as well as treatment before the tumour reaches metastatic capacity. Choroidal naevi that show no risk features should be regularly monitored and treated when tumours show growth before their diameters are over 3 mm. If the tumours are located outside the posterior pole, they should be treated with less delay. This thesis reports reference rates for tumour growth and doubling times, which could be used for earlier diagnostics. Newer imaging techniques provide new information and show new risk factors for choroidal lesions, aiming at an earlier detection.

It is likely that, in the near future, radiotherapy will remain as the principal treatment option for small CMs. However, an earlier diagnosis could possibly enable the safe use of TTT for extremely small flat tumours despite the criticism against TTT's high recurrence rates for thicker tumours. However, further studies and longer follow-ups are needed to confirm the efficacy of TTT when treating these incipient melanomas.

Metastatic disease constitutes a significant cause of mortality, including in patients with a small CM. It will be necessary, in the future, to detect and manage subclinical distant metastasis to ultimately control this malignant disease.

With earlier local treatment combined with future systemic therapies for subclinical metastasis, the improved survival of patients with UM will be accomplished. Effective local treatment modalities with effective tumour control are available for the appropriate patient selection. However, the prevention of dissemination will be the key to providing the most ideal prognosis. Currently, the most effective measure to minimize poor prognosis is the early detection of CM at a time when the tumour is small, posing the least risk for metastatic disease.

11. Conclusion

- I. An acceptable local tumour control, a low rate of side-effects other than radiation maculopathy, and the effective preservation of vision support the use of a 10-mm ruthenium¹⁰⁶ plaque in treating small CMs.
- II. 10-mm ruthenium¹⁰⁶ plaque contributes to improved visual preservation without increasing the local recurrence rate, particularly with tumours close to the fovea, and may therefore be preferable to the 15-mm plaque.
- III. Small CMs of less than 3 mm in LBD are highly unlikely to metastasize, and no known clinical risk feature can predict the development of metastases from a tumour of this size.
- IV. Small incipient CMs can be differentiated from small naevi by their growth rates, tumour doubling times, and the calculated age when the tumour originated.

12. Acknowledgements

This study was carried out at the Department of Ophthalmology, University of Helsinki and Helsinki University Central Hospital during the time period from 2011–2020. The research was supported by the Sigrid Jusélius Foundation, The Eye Foundation, the Evald ja Hilda Nissi Foundation, the Mary and Georg Ehrnrooth Foundation, the Foundation of the Eye and Tissue Bank, and the Finnish Ophthalmological Society. I am grateful for this financial support, which has enabled my research. I am grateful to the chief of the department, Jukka Moilanen, MD, for establishing ideal research facilities and for having an encouraging attitude.

My sincerest gratitude is owed to my supervisor, Professor Tero Kivelä, MD. It was summer 2010 when I first contacted Tero Kivelä to ask if I could have a summer job as a researcher aiming to complete my medical licence thesis. Tero Kivelä offered to me a project that aimed toward a doctoral thesis and degree. Possibly due to that interesting offer, I am also specialising in ophthalmology. I could not have received more suitable guidance, teaching, and support during this project. The knowledge and endless hours he provided to this project are more than I could ask for. I am thankful that he guided me through this project and taught me what high-quality research is.

I want to thank my co-writer Rana'a Al Jamal, MD for all the help and support. I further want to thank other co-writers Vappu Reijonen, PhD; Jorma Heikkonen, PhD; Sebastian Eskelin, MD; Virpi Raivio, MD; and Martin Täll, MD.

I want to thank the European Ocular Oncology Group for the opportunity to take part in the collaborative study that our group was privileged to complete. I want to thank all the members who participated in the data collection process and made this exquisite study possible.

It is a truly honour to thank Carol Shields, MD for sharing their unique data set of growing nevi that was used as reference group in the Study IV.

Warm thanks belong to Elina Rantala, MD for all the help, for all the weekends we worked together in the clinic with our own projects, and for demonstrating a hardworking attitude, which I tried to follow. I want to thank Docent Joni Turunen, MD for all the help, and for inspiration to research. I want to thank other doctoral candidates in our research group for support and, most of all, friendship, including Anni Stadigh, MD; Mari Mansikkamäki, MD; Kalle Nummi, MD; Aino Jaakkola, MD; Mamunur Rashid, MD; and Olli Ala-fossi, MD.

I am grateful to Elina Haapaniemi for her endless kindness and helpfulness.

I want to thank my thesis committee members Docent Eeva-Marja Sankila, MD and Docent Mikko Tenhunen, PhD for their support and help.

I want to thank my reviewers, Professor Kai Kaarniranta, MD, and Docent Ann Maria Paula Schalenbourg, MD, for excellent comments as well as constructive criticism.

I want to thank all my colleagues working at the Helsinki University Hospital, Department of Ophthalmology for inspiring, encouraging, and providing me with a respectful working atmosphere for residency and research.

My deepest gratitude belongs to my parents, Eira and Mika, and my brother Niko, who have supported and loved me throughout my life. I am very fortunate to have such a strong and safe background. I am sad that my beloved grandparents never saw me finishing my doctoral degree, although I know how proud they would have been.

Lastly, I want to thank the love of my life, my husband Lauri, who has encouraged me throughout this entire project by supporting, helping, and showing an example by completing his own doctoral degree a few years earlier. He inspires me each day, and it is my greatest luck to share my life with him.

13. References

1. Char DH. The management of small choroidal melanomas. *Surv Ophthalmol.* 1978;22(6):377-386.
2. Gass JD. Observation of suspected choroidal and ciliary body melanomas for evidence of growth prior to enucleation. *Ophthalmology.* 1980;87(6):523-528.
3. Augsburger JJ, Vrabec TR. Impact of delayed treatment in growing posterior uveal melanomas. *Arch Ophthalmol.* 1993;111(10):1382-1386.
4. Augsburger JJ, Corrêa ZM, Trichopoulos N. An alternative hypothesis for observed mortality rates due to metastasis after treatment of choroidal melanomas of different sizes. *Trans Am Ophthalmol Soc.* 2007;105:54-60.
5. Eskelin S, Pyrhönen S, Hahka-Kemppinen M, Tuomaala S, Kivelä T. A prognostic model and staging for metastatic uveal melanoma. *Cancer.* 2003;97(2):465-475.
6. Jampol LM, Moy CS, Murray TG, et al. The COMS randomized trial of iodine 125 brachytherapy for choroidal melanoma: IV. Local treatment failure and enucleation in the first 5 years after brachytherapy. COMS report no. 19. *Ophthalmology.* 2002;109(12):2197-2206.
7. Vrabec TR, Augsburger JJ, Gamel JW, Brady LW, Hernandez C, Woodleigh R. Impact of local tumor relapse on patient survival after cobalt 60 plaque radiotherapy. *Ophthalmology.* 1991;98(6):984-988.
8. Ophthalmic Oncology Task Force. Local recurrence significantly increases the risk of metastatic uveal melanoma. *Ophthalmology.* 2016;123:86-91.
9. Brachytherapy EZBGO. https://www.bebig.com/fileadmin/bebig_neu/user_uploads/Products/Ophthalmic_Brachytherapy/Ophthalmic_Brochure_Rev_06_English.pdf. Accessed(accessed on 28 July 2020).
10. McLaughlin CC, Wu XC, Jemal A, Martin HJ, Roche LM, Chen VW. Incidence of noncutaneous melanomas in the U.S. *Cancer.* 2005;103(5):1000-1007.
11. Damato EM, Damato BE. Detection and time to treatment of uveal melanoma in the United Kingdom: an evaluation of 2,384 patients. *Ophthalmology.* 2012;119(8):1582-1589.
12. Andreoli MT, Mieler WF, Leiderman YI. Epidemiological trends in uveal melanoma. *Br J Ophthalmol.* 2015;99(11):1550-1553.
13. Kivelä T. The epidemiological challenge of the most frequent eye cancer: retinoblastoma, an issue of birth and death. *Br J Ophthalmol.* 2009;93(9):1129-1131.
14. Singh AD, Topham A. Incidence of uveal melanoma in the United States: 1973-1997. *Ophthalmology.* 2003;110(5):956-961.
15. Singh AD, Turell ME, Topham AK. Uveal melanoma: trends in incidence, treatment, and survival. *Ophthalmology.* 2011;118(9):1881-1885.
16. Virgili G, Gatta G, Ciccolallo L, et al. Survival in patients with uveal melanoma in Europe. *Arch Ophthalmol.* 2008;126(10):1413-1418.
17. The COMS randomized trial of iodine 125 brachytherapy for choroidal melanoma: V. Twelve-year mortality rates and prognostic factors: COMS report No. 28. *Arch Ophthalmol.* 2006;124:1684-1693.

18. Diener-West M, Hawkins BS, Markowitz JA, Schachat AP. A review of mortality from choroidal melanoma. II. A meta-analysis of 5-year mortality rates following enucleation, 1966 through 1988. *Arch Ophthalmol*. 1992;110(2):245-250.
19. Diener-West M, Reynolds SM, Agugliaro DJ, et al. Development of metastatic disease after enrollment in the COMS trials for treatment of choroidal melanoma: Collaborative Ocular Melanoma Study Group Report No. 26. *Arch Ophthalmol*. 2005;123(12):1639-1643.
20. Kujala E, Makitie T, Kivela T. Very long-term prognosis of patients with malignant uveal melanoma. *Invest Ophthalmol Vis Sci*. 2003;44(11):4651-4659.
21. Gamel JW, McLean IW, McCurdy JB. Biologic distinctions between cure and time to death in 2892 patients with intraocular melanoma. *Cancer*. 1993;71(7):2299-2305.
22. Duh EJ, Schachat AP, Albert DM, Patel SM. Long-term survival in a patient with uveal melanoma and liver metastasis. *Arch Ophthalmol*. 2004;122(2):285-287.
23. Rantala ES, Hernberg M, Kivelä TT. Overall survival after treatment for metastatic uveal melanoma: a systematic review and meta-analysis. *Melanoma Research*. 2019;29(6):561-568.
24. Kujala E, Damato B, Coupland SE, et al. Staging of ciliary body and choroidal melanomas based on anatomic extent. *J Clin Oncol*. 2013;31:2825-2831.
25. Sumich P, Mitchell P, Wang JJ. Choroidal nevi in a white population: the Blue Mountains Eye Study. *Arch Ophthalmol*. 1998;116(5):645-650.
26. Singh AD, Kalyani P, Topham A. Estimating the risk of malignant transformation of a choroidal nevus. *Ophthalmology*. 2005;112(10):1784-1789.
27. Shields CL, Shields JA, Kiratli H, De Potter P, Cater JR. Risk factors for growth and metastasis of small choroidal melanocytic lesions. *Ophthalmology*. 1995;102:1351-1361.
28. Shields CL, Cater J, Shields JA, Singh AD, Santos MC, Carvalho C. Combination of clinical factors predictive of growth of small choroidal melanocytic tumors. *Arch Ophthalmol*. 2000;118(3):360-364.
29. Shields CL, Furuta M, Berman EL, et al. Choroidal nevus transformation into melanoma: analysis of 2514 consecutive cases. *Arch Ophthalmol*. 2009;127(8):981-987.
30. Damato B. Ocular treatment of choroidal melanoma in relation to the prevention of metastatic death - A personal view. *Prog Retin Eye Res*. 2018;66:187-199.
31. The C, Ocular, Melanoma, Study, Group. Factors predictive of growth and treatment of small choroidal melanoma: COMS Report No. 5. *Arch Ophthalmol*. 1997;115(12):1537-1544.
32. Singh AD, Mokashi AA, Bena JF, Jacques R, Rundle PA, Rennie IG. Small choroidal melanocytic lesions: features predictive of growth. *Ophthalmology*. 2006;113(6):1032-1039.
33. Shields CL, Dalvin LA, Ancona-Lezama D, et al. Choroidal nevus imaging features in 3,806 cases and risk factors for transformations into melanoma in 2,355 cases: The 2020 Taylor R. Smith and Victor T. Curtin Lecture. *Retina*. 2019;39(10):1840-1851.
34. Finger P, Kurli M, Shulman J. Treatment of small choroidal melanoma: An Internet Survey. *Internet J Ophthalmol Vis Sci*. 2006;5(1):1-5.
35. Augsburger JJ. Is observation really appropriate for small choroidal melanomas. *Trans Am Ophthalmol Soc*. 1993;91:147-175.
36. Robertson DM. Changing concepts in the management of choroidal melanoma. *Am J Ophthalmol*. 2003;136(1):161-170.

37. Murray TG, Sobrin L. The case for observational management of suspected small choroidal melanoma. *Arch Ophthalmol*. 2006;124(9):1342-1344.
38. Shields JA. Treating some small melanocytic choroidal lesions without waiting for growth. *Arch Ophthalmol*. 2006;124(9):1344-1346.
39. Butler P, Char DH, Zarbin M, Kroll S. Natural history of indeterminate pigmented choroidal tumors. *Ophthalmology*. 1994;101(4):710-717.
40. Augsburger JJ, Schroeder RP, Territo C, Gamel JW, Shields JA. Clinical parameters predictive of enlargement of melanocytic choroidal lesions. *Br J Ophthalmol*. 1989;73(11):911-917.
41. Desjardins L, Lumbroso L, Levy C, Plancher C, Asselain B. Risk factors for the degeneration of the choroid naevi: a retrospective study of 135 cases. *J Fr Ophtalmol*. 2001;24(6):610-616.
42. Shields CL, Shields JA. Clinical features of small choroidal melanoma. *Curr Opin Ophthalmol*. 2002;13:135-141.
43. Mashayekhi A, Siu S, Shields CL, Shields JA. Slow enlargement of choroidal nevi: a long-term follow-up study. *Ophthalmology*. 2011;118:382-388.
44. Shields CL, Shields JA. Ocular melanoma: relatively rare but requiring respect. *Clinics in dermatology*. 2009;27(1):122-133.
45. Lommatzsch PK. Results after beta-irradiation (106Ru/106Rh) of choroidal melanomas: 20 years' experience. *Br J Ophthalmol*. 1986;70(11):844-851.
46. Stallard HB. Radiotherapy for malignant melanoma of the choroid. *Br J Ophthalmol*. 1966;50(3):147-155.
47. Shields JA, Augsburger JJ, Brady LW, Day JL. Cobalt plaque therapy of posterior uveal melanomas. *Ophthalmology*. 1982;89(10):1201-1207.
48. Oosterhuis JA, Journee-de Korver HG, Kakebeeke-Kemme HM, Bleeker JC. Transpupillary thermotherapy in choroidal melanomas. *Arch Ophthalmol*. 1995;113(3):315-321.
49. Isager P, Ehlers N, Urbak SF, Overgaard J. Visual outcome, local tumour control, and eye preservation after 106Ru/Rh brachytherapy for choroidal melanoma. *Acta Oncol*. 2006;45(3):285-293.
50. Marconi DG, de Castro DG, Rebouças LM, et al. Tumor control, eye preservation, and visual outcomes of ruthenium plaque brachytherapy for choroidal melanoma. *Brachytherapy*. 2013;12(3):235-239.
51. Egan KM, Ryan LM, Gragoudas ES. Survival implications of enucleation after definitive radiotherapy for choroidal melanoma: an example of regression on time-dependent covariates. *Arch Ophthalmol*. 1998;116(3):366-370.
52. Singh AD, Kivelä T, Seregard S, Robertson D, Bena JF. Primary transpupillary thermotherapy of "small" choroidal melanoma: is it safe? *Br J Ophthalmol*. 2008;92:727-728.
53. Mashayekhi A, Shields CL, Rishi P, et al. Primary transpupillary thermotherapy for choroidal melanoma in 391 cases: importance of risk factors in tumor control. *Ophthalmology*. 2015;122:600-609.
54. De Potter P, Jamart J. Adjuvant indocyanine green in transpupillary thermotherapy for choroidal melanoma. *Ophthalmology*. 2003;110(2):406-413; discussion 413-404.

55. Harbour JW, Meredith TA, Thompson PA, Gordon ME. Transpupillary thermotherapy versus plaque radiotherapy for suspected choroidal melanomas. *Ophthalmology*. 2003;110(11):2207-2214; discussion 2215.
56. Aaberg TM, Jr., Bergstrom CS, Hickner ZJ, Lynn MJ. Long-term results of primary transpupillary thermal therapy for the treatment of choroidal malignant melanoma. *Br J Ophthalmol*. 2008;92:741-746.
57. Parrozzani R, Boccassini B, De Belvis V, Radin PP, Midena E. Long-term outcome of transpupillary thermotherapy as primary treatment of selected choroidal melanoma. *Acta Ophthalmol*. 2009;87(7):789-792.
58. Pan Y, Diddie K, Lim JI. Primary transpupillary thermotherapy for small choroidal melanomas. *Br J Ophthalmol*. 2008;92(6):747-750.
59. Win PH, Robertson DM, Buettner H, McCannel CA, Bennett SR. Extended follow-up of small melanocytic choroidal tumors treated with transpupillary thermotherapy. *Arch Ophthalmol*. 2006;124:503-506.
60. Yarovoy AA, Magaramov DA, Bulgakova ES. Which choroidal melanoma should be treated with primary transpupillary thermotherapy? Our experience from 78 patients. *Eur J Ophthalmol*. 2010;20(1):186-193.
61. Jouhi S, Jager MJ, de Geus SJR, et al. The Small Fatal Choroidal Melanoma Study. A Survey by the European Ophthalmic Oncology Group. *Am J Ophthalmol*. 2019;202:100-108.
62. Aronow ME, Topham AK, Singh AD. Uveal Melanoma: 5-Year Update on Incidence, Treatment, and Survival (SEER 1973-2013). *Ocul Oncol Pathol*. 2018;4(3):145-151.
63. Egan KM, Seddon JM, Glynn RJ, Gragoudas ES, Albert DM. Epidemiologic aspects of uveal melanoma. *Surv Ophthalmol*. 1988;32(4):239-251.
64. Espinoza G, Rosenblatt B, Harbour JW. Optical coherence tomography in the evaluation of retinal changes associated with suspicious choroidal melanocytic tumors. *Am J Ophthalmol*. 2004;137(1):90-95.
65. Kivelä T. Incidence, prevalence and epidemiology of ocular melanoma. In: *Ocular Melanoma: Advances in Diagnostic and Therapeutic Strategies*. 2014:20-38.
66. Virgili G, Gatta G, Ciccolallo L, et al. Incidence of uveal melanoma in Europe. *Ophthalmology*. 2007;114(12):2309-2315.
67. Raivio I. Uveal melanoma in Finland. An epidemiological, clinical, histological and prognostic study. *Acta Ophthalmol Suppl*. 1977(133):1-64.
68. Ghazawi FM, Darwich R, Le M, et al. Uveal melanoma incidence trends in Canada: a national comprehensive population-based study. *Br J Ophthalmol*. 2019;103(12):1872-1876.
69. Xu Y, Lou L, Wang Y, et al. Epidemiological Study of Uveal Melanoma from US Surveillance, Epidemiology, and End Results Program (2010-2015). *J Ophthalmol*. 2020;2020:3614039.
70. Park SJ, Oh CM, Kim BW, Woo SJ, Cho H, Park KH. Nationwide Incidence of Ocular Melanoma in South Korea by Using the National Cancer Registry Database (1999-2011). *Invest Ophthalmol Vis Sci*. 2015;56(8):4719-4724.
71. Vajdic CM, Krickler A, Gibling M, et al. Incidence of ocular melanoma in Australia from 1990 to 1998. *Int J Cancer*. 2003;105(1):117-122.
72. Diener-West M, Earle JD, Fine SL, et al. The COMS randomized trial of iodine 125 brachytherapy for choroidal melanoma, III: initial mortality findings. COMS Report No. 18. *Arch Ophthalmol*. 2001;119:969-982.

73. Hammer H, Oláh J, Tóth-Molnár E. Dysplastic nevi are a risk factor for uveal melanoma. *Eur J Ophthalmol*. 1996;6(4):472-474.
74. Carr S, Smith C, Wernberg J. Epidemiology and Risk Factors of Melanoma. *Surg Clin North Am*. 2020;100(1):1-12.
75. Kastelan S, Gverovic Antunica A, Beketic Oreskovic L, Salopek Rabatic J, Kasun B, Bakija I. Conjunctival Melanoma - Epidemiological Trends and Features. *Pathol Oncol Res*. 2018;24(4):787-796.
76. Shields CL, Furuta M, Thangappan A, et al. Metastasis of uveal melanoma millimeter-by-millimeter in 8033 consecutive eyes. *Arch Ophthalmol*. 2009;127(8):989-998.
77. Kivelä T. Prevalence and epidemiology of ocular melanoma. In: Murray T, Boldt HC, eds. *Ocular Melanoma: Advances in Diagnostic and Therapeutic Strategies*. London: Future Science; 2014:21-38.
78. Bergman L, Seregard S, Nilsson B, Ringborg U, Lundell G, Ragnarsson-Olding B. Incidence of uveal melanoma in Sweden from 1960 to 1998. *Invest Ophthalmol Vis Sci*. 2002;43(8):2579-2583.
79. Singh AD, Shields CL, Shields JA, Sato T. Uveal melanoma in young patients. *Arch Ophthalmol*. 2000;118(7):918-923.
80. Shields CL, Kaliki S, Arepalli S, et al. Uveal melanoma in children and teenagers. *Saudi J Ophthalmol*. 2013;27(3):197-201.
81. Al-Jamal RT, Cassoux N, Desjardins L, et al. The Pediatric Choroidal and Ciliary Body Melanoma Study: A Survey by the European Ophthalmic Oncology Group. *Ophthalmology*. 2016;123:898-907.
82. Harbour JW, Onken MD, Roberson ED, et al. Frequent mutation of BAP1 in metastasizing uveal melanomas. *Science*. 2010;330(6009):1410-1413.
83. Onken MD, Worley LA, Long MD, et al. Oncogenic mutations in GNAQ occur early in uveal melanoma. *Invest Ophthalmol Vis Sci*. 2008;49(12):5230-5234.
84. Van Raamsdonk CD, Bezrookove V, Green G, et al. Frequent somatic mutations of GNAQ in uveal melanoma and blue naevi. *Nature*. 2009;457(7229):599-602.
85. Van Raamsdonk CD, Griewank KG, Crosby MB, et al. Mutations in GNA11 in uveal melanoma. *N Engl J Med*. 2010;363(23):2191-2199.
86. Harbour JW, Roberson ED, Anbunathan H, Onken MD, Worley LA, Bowcock AM. Recurrent mutations at codon 625 of the splicing factor SF3B1 in uveal melanoma. *Nat Genet*. 2013;45(2):133-135.
87. Martin M, Masshofer L, Temming P, et al. Exome sequencing identifies recurrent somatic mutations in EIF1AX and SF3B1 in uveal melanoma with disomy 3. *Nat Genet*. 2013;45(8):933-936.
88. Bauer J, Kilic E, Vaarwater J, Bastian BC, Garbe C, de Klein A. Oncogenic GNAQ mutations are not correlated with disease-free survival in uveal melanoma. *Br J Cancer*. 2009;101(5):813-815.
89. Decatur CL, Ong E, Garg N, et al. Driver Mutations in Uveal Melanoma: Associations With Gene Expression Profile and Patient Outcomes. *JAMA Ophthalmol*. 2016;134(7):728-733.
90. Kivelä T. Uveal Malignant Melanoma: Histopathologic Features. In: Damato B, Singh A, eds. *Clinical Ophthalmic Oncology*. Vol 2. Verlag Berlin Heidelberg: Springer; 2014:113-123.

91. Damato B, Lecuona K. Conservation of eyes with choroidal melanoma by a multimodality approach to treatment: an audit of 1632 patients. *Ophthalmology*. 2004;111(5):977-983.
92. Jensen OA. Malignant melanomas of the human uvea: 25-year follow-up of cases in Denmark, 1943--1952. *Acta ophthalmologica*. 1982;60(2):161-182.
93. Aoki H, Yamada Y, Hara A, Kunisada T. Two distinct types of mouse melanocyte: differential signaling requirement for the maintenance of non-cutaneous and dermal versus epidermal melanocytes. *Development*. 2009;136(15):2511-2521.
94. Clarijs R, Schalkwijk L, Ruiter DJ, de Waal RM. Lack of lymphangiogenesis despite coexpression of VEGF-C and its receptor Flt-4 in uveal melanoma. *Invest Ophthalmol Vis Sci*. 2001;42(7):1422-1428.
95. Holly EA, Aston DA, Char DH, Kristiansen JJ, Ahn DK. Uveal melanoma in relation to ultraviolet light exposure and host factors. *Cancer Res*. 1990;50(18):5773-5777.
96. Mallet JD, Gendron SP, Drigeard Desgarnier MC, Rochette PJ. Implication of ultraviolet light in the etiology of uveal melanoma: A review. *Photochem Photobiol*. 2014;90(1):15-21.
97. Singh AD, Rennie IG, Seregard S, Giblin M, McKenzie J. Sunlight exposure and pathogenesis of uveal melanoma. *Surv Ophthalmol*. 2004;49(4):419-428.
98. Gandini S, Sera F, Cattaruzza MS, et al. Meta-analysis of risk factors for cutaneous melanoma: II. Sun exposure. *Eur J Cancer*. 2005;41(1):45-60.
99. Shah CP, Weis E, Lajous M, Shields JA, Shields CL. Intermittent and chronic ultraviolet light exposure and uveal melanoma: a meta-analysis. *Ophthalmology*. 2005;112(9):1599-1607.
100. Gu  nel P, Laforest L, Cyr D, et al. Occupational risk factors, ultraviolet radiation, and ocular melanoma: a case-control study in France. *Cancer Causes Control*. 2001;12(5):451-459.
101. Vidal JL, Bacin F, Albuisson E, et al. ["Melanoma 92". epidemiological study of uveal melanoma in France]. *J Fr Ophtalmol*. 1995;18(8-9):520-528.
102. Durante MA, Field MG, Sanchez MI, et al. Genomic evolution of uveal melanoma arising in ocular melanocytosis. *Cold Spring Harb Mol Case Stud*. 2019;5(4).
103. Shields CL, Say EAT, Hasanreisoglu M, et al. Cytogenetic abnormalities in uveal melanoma based on tumor features and size in 1059 patients: the 2016 W. Richard Green lecture. *Ophthalmology*. 2017;124:609-618.
104. Gonder JR, Shields JA, Albert DM, Augsburger JJ, Lavin PT. Uveal malignant melanoma associated with ocular and oculodermal melanocytosis. *Ophthalmology*. 1982;89(8):953-960.
105. Shields CL, Furuta M, Mashayekhi A, et al. Clinical spectrum of choroidal nevi based on age at presentation in 3422 consecutive eyes. *Ophthalmology*. 2008;115(3):546-552.e542.
106. Chien JL, Sioufi K, Surakiatchanukul T, Shields JA, Shields CL. Choroidal nevus: a review of prevalence, features, genetics, risks, and outcomes. *Curr Opin Ophthalmol*. 2017;28(3):228-237.
107. Kivela T, Eskelin S. Transformation of nevus to melanoma. *Ophthalmology*. 2006;113(5):887-888.e881.
108. Eskelin S, Kivela T. Mode of presentation and time to treatment of uveal melanoma in Finland. *Br J Ophthalmol*. 2002;86(3):333-338.
109. Naumann G, Yanoff M, Zimmerman LE. Histogenesis of malignant melanomas of the uvea. I. Histopathologic characteristics of nevi of the choroid and ciliary body. *Arch Ophthalmol*. 1966;76(6):784-796.

110. Yanoff M, Zimmerman LE. Histogenesis of malignant melanomas of the uvea. 3. The relationship of congenital ocular melanocytosis and neurofibromatosis in uveal melanomas. *Arch Ophthalmol*. 1967;77(3):331-336.
111. Qiu M, Shields CL. Choroidal Nevus in the United States Adult Population: Racial Disparities and Associated Factors in the National Health and Nutrition Examination Survey. *Ophthalmology*. 2015;122(10):2071-2083.
112. Jonas JB, You QS, Xu L, Wang YX. Choroidal nevi in adult chinese. *Ophthalmology*. 2008;115(6):1102-1102.e1101.
113. Ganley JP, Comstock GW. Benign nevi and malignant melanomas of the choroid. *Am J Ophthalmol*. 1973;76(1):19-25.
114. Abdel-Rahman MH, Pilarski R, Cebulla CM, et al. Germline BAP1 mutation predisposes to uveal melanoma, lung adenocarcinoma, meningioma, and other cancers. *J Med Genet*. 2011;48:856-859.
115. Wiesner T, Obenaus AC, Murali R, et al. Germline mutations in BAP1 predispose to melanocytic tumors. *Nat Genet*. 2011;43(10):1018-1021.
116. Testa JR, Cheung M, Pei J, et al. Germline BAP1 mutations predispose to malignant mesothelioma. *Nat Genet*. 2011;43(10):1022-1025.
117. Njauw CN, Kim I, Piris A, et al. Germline BAP1 inactivation is preferentially associated with metastatic ocular melanoma and cutaneous-ocular melanoma families. *PLoS One*. 2012;7(4):e35295.
118. Carbone M, Ferris LK, Baumann F, et al. BAP1 cancer syndrome: malignant mesothelioma, uveal and cutaneous melanoma, and MBAITs. *J Transl Med*. 2012;10:179.
119. Singh AD, Shields CL, De Potter P, et al. Familial Uveal Melanoma: Clinical Observations on 56 Patients. *Arch Ophthalmol*. 1996;114(4):392-399.
120. Barker-Griffith AE, Streeten BW. Familial uveal melanoma: a report of four cases in two families and literature review. *Can J Ophthalmol*. 2004;39(4):403-408.
121. Field MG, Harbour JW. Recent developments in prognostic and predictive testing in uveal melanoma. *Curr Opin Ophthalmol*. 2014;25(3):234-239.
122. Gupta MP, Lane AM, DeAngelis MM, et al. Clinical Characteristics of Uveal Melanoma in Patients With Germline BAP1 Mutations. *JAMA Ophthalmol*. 2015;133(8):881-887.
123. Aoude LG, Vajdic CM, Krickler A, Armstrong B, Hayward NK. Prevalence of germline BAP1 mutation in a population-based sample of uveal melanoma cases. *Pigment Cell Melanoma Res*. 2013;26(2):278-279.
124. Turunen JA, Markkinen S, Wilska R, et al. BAP1 Germline Mutations in Finnish Patients with Uveal Melanoma. *Ophthalmology*. 2016;123(5):1112-1117.
125. Repo P, Jäntti JE, Järvinen RS, et al. Germline loss-of-function variants in MBD4 are rare in Finnish patients with uveal melanoma. *Pigment Cell Melanoma Res*. 2020.
126. Lains I, Bartosch C, Mondim V, et al. Second Primary Neoplasms in Patients With Uveal Melanoma: A SEER Database Analysis. *Am J Ophthalmol*. 2016;165:54-64.
127. Derrien AC, Rodrigues M, Eeckhoutte A, et al. Germline MBD4 mutations and predisposition to uveal melanoma. *J Natl Cancer Inst*. 2020.
128. Johansson PA, Stark A, Palmer JM, et al. Prolonged stable disease in a uveal melanoma patient with germline MBD4 nonsense mutation treated with pembrolizumab and ipilimumab. *Immunogenetics*. 2019;71(5-6):433-436.

129. Rodrigues M, Mobuchon L, Houy A, et al. Outlier response to anti-PD1 in uveal melanoma reveals germline MBD4 mutations in hypermutated tumors. *Nat Commun.* 2018;9(1):1866.
130. Hodi FS, Chiarion-Sileni V, Gonzalez R, et al. Nivolumab plus ipilimumab or nivolumab alone versus ipilimumab alone in advanced melanoma (CheckMate 067): 4-year outcomes of a multicentre, randomised, phase 3 trial. *Lancet Oncol.* 2018;19(11):1480-1492.
131. Schachter J, Ribas A, Long GV, et al. Pembrolizumab versus ipilimumab for advanced melanoma: final overall survival results of a multicentre, randomised, open-label phase 3 study (KEYNOTE-006). *Lancet.* 2017;390(10105):1853-1862.
132. Schank TE, Hassel JC. Immunotherapies for the Treatment of Uveal Melanoma-History and Future. *Cancers.* 2019;11(8).
133. Kivelä T. Diagnosis of uveal melanoma. *Dev Ophthalmol.* 2012;49:1-15.
134. Schalenbourg A, Zografos L. Clinical Ophthalmic Oncology. In: Damato B, Singh A, eds. *Uveal tumors.* 2 ed. Verlag Berlin Heidelberg: Springer; 2014:93-98.
135. Papadopoulou D, Moulin AP, Zografos L, Schalenbourg A. How would you manage this small melanocytic choroidal tumour? Small choroidal melanoma hiding a large extrascleral extension. *Br J Ophthalmol.* 2012;96(12):1530, 1539-1540.
136. Sambuelli R, Luna JD, Reviglio VE, Aoki A, Juarez CP. Small choroidal melanoma with massive extraocular extension: invasion through posterior scleral emissary channels. *Int Ophthalmol.* 2001;24(4):213-218.
137. Duffin RM, Straatsma BR, Foos RY, Kerman BM. Small malignant melanoma of the choroid with extraocular extension. *Arch Ophthalmol.* 1981;99(10):1827-1830.
138. Lemaître S, Zmuda M, Jacomet PV, et al. Small Choroidal Melanoma Revealed by a Large Extrascleral Extension. *Ocul Oncol Pathol.* 2017;3(3):240-246.
139. Shields CL, Santos MC, Shields JA, Singh AD, Eagle RC, Jr. Extraocular extension of unrecognized choroidal melanoma simulating a primary optic nerve tumor: report of two cases. *Ophthalmology.* 1999;106(7):1349-1352.
140. Coupland SE, Campbell I, Damato B. Routes of extraocular extension of uveal melanoma: risk factors and influence on survival probability. *Ophthalmology.* 2008;115(10):1778-1785.
141. van Beek JG, Koopmans AE, Vaarwater J, et al. The prognostic value of extraocular extension in relation to monosomy 3 and gain of chromosome 8q in uveal melanoma. *Invest Ophthalmol Vis Sci.* 2014;55(3):1284-1291.
142. McLean IW, Foster WD, Zimmerman LE, Martin DG. Inferred natural history of uveal melanoma. *Invest Ophthalmol Vis Sci.* 1980;19(7):760-770.
143. The Collaborative Ocular Melanoma Study Group. Mortality in patients with small choroidal melanoma. COMS report no. 4. . *Arch Ophthalmol.* 1997;115(7):886-893.
144. Augsburger JJ, Correa ZM, Trichopoulos N, Shaikh A. Size overlap between benign melanocytic choroidal nevi and choroidal malignant melanomas. *Invest Ophthalmol Vis Sci.* 2008;49(7):2823-2828.
145. Kivelä T, Simpson R, Grossniklaus HE, et al. Uveal Melanoma. In: Mahul BA, ed. *AJCC Cancer Staging Manual.* 8 ed. Chicago: Springer; 2017:805-817.
146. Finger PT. The 7th edition AJCC staging system for eye cancer: an international language for ophthalmic oncology. *Arch Pathol Lab Med.* 2009;133(8):1197-1198.

147. Gass JD. Problems in the differential diagnosis of choroidal nevi and malignant melanoma. XXXIII Edward Jackson Memorial lecture. *Transactions Section on Ophthalmology American Academy of Ophthalmology and Otolaryngology*. 1977;83(1):19-48.
148. Shields CL, Demirci H, Materin MA, Marr BP, Mashayekhi A, Shields JA. Clinical factors in the identification of small choroidal melanoma. *Can J Ophthalmol*. 2004;39(4):351-357.
149. Harbour JW, Paez-Escamilla M, Cai L, Walter SD, Augsburger JJ, Correa ZM. Are Risk Factors for Growth of Choroidal Nevi Associated With Malignant Transformation? Assessment With a Validated Genomic Biomarker. *Am J Ophthalmol*. 2019;197:168-179.
150. Shields CL, Dalvin LA, Yu MD, et al. Choroidal nevus transformation into melanoma per millimeter increment in thickness using multimodal imaging in 2355 cases: The 2019 Wendell L. Hughes Lecture. *Retina*. 2019;39(10):1852-1860.
151. Kivelä T, Eskelin S, Mäkitie T, Summanen P. Exudative retinal detachment from malignant uveal melanoma: predictors and prognostic significance. *Invest Ophthalmol Vis Sci*. 2001;42(9):2085-2093.
152. Yu MD, Dalvin LA, Ancona-Lezama D, et al. Choriocapillaris Compression Correlates with Choroidal Nevus-Associated Subretinal Fluid: OCT Analysis of 3431 Cases. *Ophthalmology*. 2020.
153. Gonder JR, Augsburger JJ, McCarthy EF, Shields JA. Visual loss associated with choroidal nevi. *Ophthalmology*. 1982;89(8):961-965.
154. Shields CL, Mashayekhi A, Materin MA, et al. Optical coherence tomography of choroidal nevus in 120 patients. *Retina*. 2005;25(3):243-252.
155. Muscat S, Parks S, Kemp E, Keating D. Secondary retinal changes associated with choroidal naevi and melanomas documented by optical coherence tomography. *Br J Ophthalmol*. 2004;88(1):120-124.
156. Davidorf FH, Lang JR. The natural history of malignant melanoma of the choroid: small vs large tumors. *Transactions Section on Ophthalmology American Academy of Ophthalmology and Otolaryngology*. 1975;79(2):310-320.
157. Bove R, Char DH. Nondiagnosed uveal melanomas. *Ophthalmology*. 2004;111(3):554-557.
158. Shields JA, Rodrigues MM, Sarin LK, Tasman WS, Annesley WH, Jr. Lipofuscin pigment over benign and malignant choroidal tumors. *Transactions Section on Ophthalmology American Academy of Ophthalmology and Otolaryngology*. 1976;81(5):871-881.
159. Shields CL, Bianciotto C, Pirondini C, Materin MA, Harmon SA, Shields JA. Autofluorescence of orange pigment overlying small choroidal melanoma. *Retina*. 2007;27(8):1107-1111.
160. Boldt HC, Byrne SF, Gilson MM, et al. Baseline echographic characteristics of tumors in eyes of patients enrolled in the Collaborative Ocular Melanoma Study: COMS report no. 29. *Ophthalmology*. 2008;115(8):1390-1397, 1397.e1391-1392.
161. Bedi DG, Gombos DS, Ng CS, Singh S. Sonography of the eye. *AJR Am J Roentgenol*. 2006;187(4):1061-1072.
162. Fuller DG, Snyder WB, Hutton WL, Vaiser A. Ultrasonographic features of choroidal malignant melanomas. *Arch Ophthalmol*. 1979;97(8):1465-1472.
163. Goldberg MF, Hodes BL. Ultrasonographic diagnosis of choroidal malignant melanoma. *Surv Ophthalmol*. 1977;22(1):29-40.
164. Hasenfratz G. Acute vitreous hemorrhage - possibilities for differential diagnostic, echographic assessment. *Fortschr Ophthalmol*. 1990;87(6):641-645.

165. Jacobsen BH, Ricks C, Harrie RP. Ocular ultrasound versus MRI in the detection of extrascleral extension in a patient with choroidal melanoma. *BMC Ophthalmol.* 2018;18(1):320.
166. Histopathologic characteristics of uveal melanomas in eyes enucleated from the Collaborative Ocular Melanoma Study. COMS report no. 6. *Am J Ophthalmol.* 1998;125:745-766.
167. Shields CL, Maktabi AM, Jahnle E, Mashayekhi A, Lally SE, Shields JA. Halo nevus of the choroid in 150 patients: the 2010 Henry van Dyke Lecture. *Arch Ophthalmol.* 2010;128(7):859-864.
168. Davila JR, Mruthyunjaya P. Updates in imaging in ocular oncology. *F1000Res.* 2019;8:F1000 Faculty Rev-1706.
169. Toledo JJ, Asencio M, García JR, Morales LA, Tomkinson C, Cajigal C. OCT Angiography: Imaging of Choroidal and Retinal Tumors. *Ophthalmol Retina.* 2018;2(6):613-622.
170. McNabb RP, Grewal DS, Mehta R, et al. Wide field of view swept-source optical coherence tomography for peripheral retinal disease. *Br J Ophthalmol.* 2016;100(10):1377-1382.
171. Grewal DS, Bhullar PK, Pasricha ND, et al. Intraoperative 4-Dimensional Microscope-Integrated Optical Coherence Tomography-Guided 27-Gauge Transvitreal Choroidal Biopsy for Choroidal Melanoma. *Retina.* 2017;37(4):796-799.
172. Potsaid B, Baumann B, Huang D, et al. Ultrahigh speed 1050nm swept source/Fourier domain OCT retinal and anterior segment imaging at 100,000 to 400,000 axial scans per second. *Opt Express.* 2010;18(19):20029-20048.
173. Weill Y, Brosh K, Levi Vineberg T, et al. Enhanced depth imaging in swept-source optical coherence tomography: Improving visibility of choroid and sclera, a masked study. *Eur J Ophthalmol.* 2019;1120672119863560.
174. Mrejen S, Spaide RF. Optical coherence tomography: imaging of the choroid and beyond. *Surv Ophthalmol.* 2013;58(5):387-429.
175. Daitch Z, Shields CL, Say EA, Mashayekhi A, Shields JA. Submillimeter choroidal melanoma detection by enhanced depth imaging optical coherence tomography in a patient with oculodermal melanocytosis. *Retin Cases Brief Rep.* 2016;10(1):6-10.
176. Jonna G, Daniels AB. Enhanced Depth Imaging OCT of Ultrasonographically Flat Choroidal Nevi Demonstrates 5 Distinct Patterns. *Ophthalmol Retina.* 2019;3(3):270-277.
177. Shields CL, Kaliki S, Rojanaporn D, Ferenczy SR, Shields JA. Enhanced depth imaging optical coherence tomography of small choroidal melanoma: comparison with choroidal nevus. *Arch Ophthalmol.* 2012;130(7):850-856.
178. Rovati L, Docchio F. Autofluorescence methods in ophthalmology. *J Biomed Opt.* 2004;9(1):9-21.
179. Eldred GE, Katz ML. Fluorophores of the human retinal pigment epithelium: separation and spectral characterization. *Exp Eye Res.* 1988;47(1):71-86.
180. Cennamo G, Romano MR, Velotti N, Breve MA, de Crecchio G, Cennamo G. Fundus autofluorescence of choroidal nevi and melanoma. *Acta Ophthalmol.* 2018;96(1):e102-e104.
181. Gunduz K, Pulido JS, Ezzat K, Salomao D, Hann C. Review of fundus autofluorescence in choroidal melanocytic lesions. *Eye (Lond).* 2009;23(3):497-503.
182. Gündüz K, Pulido JS, Pulido JE, Link T. Correlation of fundus autofluorescence with fluorescein and indocyanine green angiography in choroidal melanocytic lesions. *Retina.* 2008;28(9):1257-1264.

183. Motiani MV, McCannel CA, Almanzor R, McCannel TA. Diagnosis of Choroidal Melanoma in Dense Asteroid Hyalosis. *Semin Ophthalmol*. 2017;32(2):257-259.
184. Pellegrini M, Corvi F, Invernizzi A, Ravera V, Cereda MG, Staurengi G. Swept-source optical coherence tomography angiography in choroidal melanoma: An analysis of 22 consecutive cases. *Retina*. 2019;39(8):1510-1519.
185. Garcia-Arumi Fuste C, Peralta Iturburu F, Garcia-Arumi J. Is optical coherence tomography angiography helpful in the differential diagnosis of choroidal nevus versus melanoma? *European journal of ophthalmology*. 2019:Epub ahead of print.
186. Mantel I, Uffer S, Zografos L. Peripheral exudative hemorrhagic chorioretinopathy: a clinical, angiographic, and histologic study. *Am J Ophthalmol*. 2009;148(6):932-938.e931.
187. Flindall RJ, Gass JD. A histopathologic fluorescein angiographic correlative study of malignant melanomas of the choroid. *Can J Ophthalmol*. 1971;6(4):258-267.
188. Hayreh SS. Choroidal melanomata. Fluorescence angiographic and histopathological study. *Br J Ophthalmol*. 1970;54(3):145-160.
189. Pettit TH, Barton A, Foos RY, Christensen RE. Fluorescein angiography of choroidal melanomas. *Arch Ophthalmol*. 1970;83(1):27-38.
190. Eskelin S, Pyrhönen S, Summanen P, Hahka-Kemppinen M, Kivelä T. Tumor doubling times in metastatic malignant melanoma of the uvea: tumor progression before and after treatment. *Ophthalmology*. 2000;107(8):1443-1449.
191. Char DH, Kroll S, Phillips TL. Uveal melanoma. Growth rate and prognosis. *Arch Ophthalmol*. 1997;115:1014-1018.
192. Augsburger JJ, Gonder JR, Amsel J, Shields JA, Donoso LA. Growth rates and doubling times of posterior uveal melanomas. *Ophthalmology*. 1984;91:1709-1715.
193. Eskelin S, Kivelä T. Author's reply. *Ophthalmology*. 2001;108(5):830-831.
194. You QS, Xu L, Jonas JB, Wang S, Yang H. Change in choroidal nevi during a 5-year follow-up study: the Beijing Eye Study. *Br J Ophthalmol*. 2010;94(5):575-578.
195. Shields CL, Lim LS, Dalvin LA, Shields JA. Small choroidal melanoma: detection with multimodal imaging and management with plaque radiotherapy or AU-011 nanoparticle therapy. *Curr Opin Ophthalmol*. 2019;30(3):206-214.
196. Sahel JA, Pesavento R, Frederick AR, Jr., Albert DM. Melanoma arising de novo over a 16-month period. *Arch Ophthalmol*. 1988;106:381-385.
197. Aleksidze N, Medina CA, Singh AD. De novo evolution of a small choroidal melanoma. *Ocul Oncol Pathol*. 2015;1:83-87.
198. Kivelä T. Observed Growth in Small Melanocytic Choroidal Tumors. *Arch Ophthalmol*. 2006;124(4):607-608.
199. Shields CL, Materin MA, Teixeira L, Mashayekhi A, Ganguly A, Shields JA. Small choroidal melanoma with chromosome 3 monosomy on fine-needle aspiration biopsy. *Ophthalmology*. 2007;114(10):1919-1924.
200. Eide N, Walaas L. Fine-needle aspiration biopsy and other biopsies in suspected intraocular malignant disease: a review. *Acta Ophthalmol*. 2009;87(6):588-601.
201. Bagger MM. Intraocular biopsy of uveal melanoma Risk assessment and identification of genetic prognostic markers. *Acta Ophthalmol*. 2018;96 Suppl A112:1-28.
202. Seider MI, Mruthyunjaya P. Molecular prognostics for uveal melanoma. *Retina*. 2018;38(2):211-219.

203. Onken MD, Worley LA, Char DH, et al. Collaborative Ocular Oncology Group report number 1: prospective validation of a multi-gene prognostic assay in uveal melanoma. *Ophthalmology*. 2012;119(8):1596-1603.
204. Midena E, Segato T, Piermarocchi S, Boccato P. Fine needle aspiration biopsy in ophthalmology. *Surv Ophthalmol*. 1985;29(6):410-422.
205. Midena E, Bonaldi L, Parrozzani R, Tebaldi E, Boccassini B, Vujosevic S. In vivo detection of monosomy 3 in eyes with medium-sized uveal melanoma using transscleral fine needle aspiration biopsy. *Eur J Ophthalmol*. 2006;16(3):422-425.
206. Accuracy of diagnosis of choroidal melanomas in the Collaborative Ocular Melanoma Study. COMS report no. 1. *Arch Ophthalmol*. 1990;108:1268-1273.
207. The Collaborative Ocular Melanoma Study Group. Accuracy of Diagnosis of Choroidal Melanomas in the Collaborative Ocular Melanoma Study: COMS Report No. 1. *Arch Ophthalmol*. 1990;108(9):1268-1273.
208. Augsburger JJ, Correa ZM, Schneider S, et al. Diagnostic transvitreal fine-needle aspiration biopsy of small melanocytic choroidal tumors in nevus versus melanoma category. *Trans Am Ophthalmol Soc*. 2002;100:225-232; discussion 232-224.
209. Corrêa ZM, Augsburger JJ. Indications for fine needle aspiration biopsy of posterior segment intraocular tumors. *Am J Ophthalmol*. 2019;207:45-61.
210. Angi M, Kalirai H, Taktak A, et al. Prognostic biopsy of choroidal melanoma: an optimised surgical and laboratory approach. *Br J Ophthalmol*. 2017;101:1143-1146.
211. Shields CL, Say EAT, Hasanreisoglu M, et al. Personalized prognosis of uveal melanoma based on cytogenetic profile in 1059 patients over an 8-year period: the 2017 Harry S. Gradle lecture. *Ophthalmology*. 2017;124:1523-1531.
212. Nagiel A, McCannel CA, Moreno C, McCannel TA. Vitrectomy-assisted biopsy for molecular prognostication of choroidal melanoma 2 mm or less in thickness with a 27-gauge cutter. *Retina*. 2017;37:1377-1382.
213. Bagger M, Tebering JF, Kiilgaard JF. The ocular consequences and applicability of minimally invasive 25-gauge transvitreal retinochoroidal biopsy. *Ophthalmology*. 2013;120(12):2565-2572.
214. Vichitvejpaisal P, Dalvin LA, Mazloumi M, Ewens KG, Ganguly A, Shields CL. Genetic analysis of uveal melanoma in 658 patients using the Cancer Genome Atlas classification of uveal melanoma as A, B, C, and D. *Ophthalmology*. 2019;126:1445-1453.
215. Bagger M, Andersen MT, Heegaard S, Andersen MK, Kiilgaard JF. Transvitreal retinochoroidal biopsy provides a representative sample from choroidal melanoma for detection of chromosome 3 aberrations. *Invest Ophthalmol Vis Sci*. 2015;56:5917-5924.
216. McCannel TA, Chang MY, Burgess BL. Multi-year follow-up of fine-needle aspiration biopsy in choroidal melanoma. *Ophthalmology*. 2012;119(3):606-610.
217. Chang MY, McCannel TA. Comparison of uveal melanoma cytopathologic sample retrieval in trans-scleral versus vitrectomy-assisted transvitreal fine needle aspiration biopsy. *Br J Ophthalmol*. 2014;98(12):1654-1658.
218. Klufas MA, Itty S, McCannel CA, Glasgow BJ, Moreno C, McCannel TA. Variable Results for Uveal Melanoma-Specific Gene Expression Profile Prognostic Test in Choroidal Metastasis. *JAMA Ophthalmol*. 2015;133(9):1073-1076.

219. Augsburger JJ, Corrêa ZM, Augsburger BD. Frequency and implications of discordant gene expression profile class in posterior uveal melanomas sampled by fine needle aspiration biopsy. *Am J Ophthalmol*. 2015;159(2):248-256.
220. Freton A, Chin KJ, Raut R, Tena LB, Kivelä T, Finger PT. Initial PET/CT staging for choroidal melanoma: AJCC correlation and second nonocular primaries in 333 patients. *Eur J Ophthalmol*. 2012;22(2):236-243.
221. Reznitsky F, Klausen S, Donia M, Scheie D. Rare cause of spontaneous haemothorax: mediastinal and distant lymph node metastases from uveal melanoma. *BMJ Case Rep*. 2019;12(11).
222. Dithmar S, Diaz CE, Grossniklaus HE. Intraocular melanoma spread to regional lymph nodes: report of two cases. *Retina*. 2000;20(1):76-79.
223. AJCC Ophthalmic Oncology Task Force. International validation of the American Joint Committee on Cancer's 7th edition classification of uveal melanoma. *JAMA Ophthalmol*. 2015;133:376-383.
224. McLean IW, Foster WD, Zimmerman LE, Gamel JW. Modifications of Callender's classification of uveal melanoma at the Armed Forces Institute of Pathology. *Am J Ophthalmol*. 1983;96(4):502-509.
225. Hamadeh F, Medina CA, Singh AD, Reynolds JP, Biscotti CV. Uveal melanoma: An analysis of cellular features and comparison to monosomy 3 status. *Diagn Cytopathol*. 2016;44(5):377-383.
226. McLean IW, Foster WD, Zimmerman LE. Uveal melanoma: location, size, cell type, and enucleation as risk factors in metastasis. *Hum Pathol*. 1982;13(2):123-132.
227. Seddon JM, Albert DM, Lavin PT, Robinson N. A prognostic factor study of disease-free interval and survival following enucleation for uveal melanoma. *Arch Ophthalmol*. 1983;101(12):1894-1899.
228. Seregard S, Kock E. Prognostic indicators following enucleation for posterior uveal melanoma. A multivariate analysis of long-term survival with minimized loss to follow-up. *Acta Ophthalmol Scand*. 1995;73(4):340-344.
229. Gamel JW, McLean IW, Foster WD, Zimmerman LE. Uveal melanomas: correlation of cytologic features with prognosis. *Cancer*. 1978;41(5):1897-1901.
230. Singh AD, Shields CL, Shields JA. Prognostic factors in uveal melanoma. *Melanoma Res*. 2001;11(3):255-263.
231. Paul E, Parnell B, M F. Prognosis of malignant melanomas of the choroid and ciliary body. *Int Ophthalmol Clin*. 1962;2:387-402.
232. Al-Jamal RT, Makitie T, Kivela T. Nucleolar diameter and microvascular factors as independent predictors of mortality from malignant melanoma of the choroid and ciliary body. *Invest Ophthalmol Vis Sci*. 2003;44:2381-2389.
233. Sorensen FB, Gamel JW, McCurdy J. Stereologic estimation of nucleolar volume in ocular melanoma: a comparative study of size estimators with prognostic impact. *Hum Pathol*. 1993;24(5):513-518.
234. McLean IW, Sibug ME, Becker RL, McCurdy JB. Uveal melanoma: the importance of large nucleoli in predicting patient outcome--an automated image analysis study. *Cancer*. 1997;79(5):982-988.

235. Mäkitie T, Summanen P, Tarkkanen A, Kivelä T. Microvascular density in predicting survival of patients with choroidal and ciliary body melanoma. *Invest Ophthalmol Vis Sci.* 1999;40(11):2471-2480.
236. Bronkhorst IH, Jager MJ. Uveal melanoma: the inflammatory microenvironment. *J Innate Immun.* 2012;4(5-6):454-462.
237. Mäkitie T, Summanen P, Tarkkanen A, Kivelä T. Tumor-infiltrating macrophages (CD68(+)) cells and prognosis in malignant uveal melanoma. *Invest Ophthalmol Vis Sci.* 2001;42(7):1414-1421.
238. Toivonen P, Mäkitie T, Kujala E, Kivela T. Macrophages and microcirculation in regressed and partially regressed irradiated choroidal and ciliary body melanomas. *Curr Eye Res.* 2003;27(4):237-245.
239. All-Ericsson C, Girnita L, Seregard S, Bartolazzi A, Jager MJ, Larsson O. Insulin-like growth factor-1 receptor in uveal melanoma: a predictor for metastatic disease and a potential therapeutic target. *Invest Ophthalmol Vis Sci.* 2002;43(1):1-8.
240. Jager MJ, Hurks HM, Levitskaya J, Kiessling R. HLA expression in uveal melanoma: there is no rule without some exception. *Hum Immunol.* 2002;63(6):444-451.
241. Folberg R, Pe'er J, Gruman LM, et al. The morphologic characteristics of tumor blood vessels as a marker of tumor progression in primary human uveal melanoma: a matched case-control study. *Hum Pathol.* 1992;23(11):1298-1305.
242. Mäkitie T, Summanen P, Tarkkanen A, Kivelä T. Microvascular loops and networks as prognostic indicators in choroidal and ciliary body melanomas. *J Natl Cancer Inst.* 1999;91(4):359-367.
243. Kivelä T, Mäkitie T, Al-Jamal RT, Toivonen P. Microvascular loops and networks in uveal melanoma. *Can J Ophthalmol.* 2004;39(4):409-421.
244. Bronkhorst IH, Jager MJ. Inflammation in uveal melanoma. *Eye.* 2013;27(2):217-223.
245. Damato B, Duke C, Coupland SE, et al. Cytogenetics of uveal melanoma: a 7-year clinical experience. *Ophthalmology.* 2007;114(10):1925-1931.
246. McLean MJ, Foster WD, Zimmerman LE. Prognostic factors in small malignant melanomas of choroid and ciliary body. *Arch Ophthalmol.* 1977;95(1):48-58.
247. Karlsson M, Boeryd B, Carstensen J, et al. Correlations of Ki-67 and PCNA to DNA ploidy, S-phase fraction and survival in uveal melanoma. *Eur J Cancer.* 1996;32a(2):357-362.
248. An J, Wan H, Zhou X, et al. A comparative transcriptomic analysis of uveal melanoma and normal uveal melanocyte. *PLoS One.* 2011;6(1):e16516.
249. Brantley MA, Jr., Harbour JW. Deregulation of the Rb and p53 pathways in uveal melanoma. *Am J Pathol.* 2000;157(6):1795-1801.
250. McCannel TA, Burgess BL, Rao NP, Nelson SF, Straatsma BR. Identification of Candidate Tumor Oncogenes by Integrative Molecular Analysis of Choroidal Melanoma Fine-Needle Aspiration Biopsy Specimens. *Arch Ophthalmol.* 2010;128(9):1170-1177.
251. Prescher G, Bornfeld N, Friedrichs W, Seeber S, Becher R. Cytogenetics of twelve cases of uveal melanoma and patterns of nonrandom anomalies and isochromosome formation. *Cancer Genet Cytogenet.* 1995;80(1):40-46.
252. Sisley K, Rennie IG, Parsons MA, et al. Abnormalities of chromosomes 3 and 8 in posterior uveal melanoma correlate with prognosis. *Genes Chromosomes Cancer.* 1997;19(1):22-28.

253. Tschentscher F, Husing J, Holter T, et al. Tumor classification based on gene expression profiling shows that uveal melanomas with and without monosomy 3 represent two distinct entities. *Cancer Res.* 2003;63(10):2578-2584.
254. Onken MD, Worley LA, Ehlers JP, Harbour JW. Gene expression profiling in uveal melanoma reveals two molecular classes and predicts metastatic death. *Cancer Res.* 2004;64(20):7205-7209.
255. Prescher G, Bornfeld N, Horsthemke B, Becher R. Chromosomal aberrations defining uveal melanoma of poor prognosis. *Lancet.* 1992;339(8794):691-692.
256. Prescher G, Bornfeld N, Hirche H, Horsthemke B, Jöckel KH, Becher R. Prognostic implications of monosomy 3 in uveal melanoma. *Lancet.* 1996;347(9010):1222-1225.
257. Caines R, Eleuteri A, Kalirai H, et al. Cluster analysis of multiplex ligation-dependent probe amplification data in choroidal melanoma. *Mol Vis.* 2015;21:1-11.
258. Cassoux N, Rodrigues MJ, Plancher C, et al. Genome-wide profiling is a clinically relevant and affordable prognostic test in posterior uveal melanoma. *Br J Ophthalmol.* 2014;98(6):769-774.
259. Sisley K, Rennie IG, Cottam DW, Potter AM, Potter CW, Rees RC. Cytogenetic findings in six posterior uveal melanomas: involvement of chromosomes 3, 6, and 8. *Genes Chromosomes Cancer.* 1990;2(3):205-209.
260. Höglund M, Gisselsson D, Hansen GB, et al. Dissecting karyotypic patterns in malignant melanomas: temporal clustering of losses and gains in melanoma karyotypic evolution. *Int J Cancer.* 2004;108(1):57-65.
261. Ehlers JP, Worley L, Onken MD, Harbour JW. Integrative genomic analysis of aneuploidy in uveal melanoma. *Clin Cancer Res.* 2008;14(1):115-122.
262. Damato B, Dopierala JA, Coupland SE. Genotypic profiling of 452 choroidal melanomas with multiplex ligation-dependent probe amplification. *Clin Cancer Res.* 2010;16(24):6083-6092.
263. Trolet J, Hupé P, Huon I, et al. Genomic profiling and identification of high-risk uveal melanoma by array CGH analysis of primary tumors and liver metastases. *Invest Ophthalmol Vis Sci.* 2009;50(6):2572-2580.
264. White VA, Chambers JD, Courtright PD, Chang WY, Horsman DE. Correlation of cytogenetic abnormalities with the outcome of patients with uveal melanoma. *Cancer.* 1998;83(2):354-359.
265. Dogrusöz M, Kroes WG, van Duinen SG, et al. Radiation Treatment Affects Chromosome Testing in Uveal Melanoma. *Invest Ophthalmol Vis Sci.* 2015;56(10):5956-5964.
266. Field MG, Durante MA, Anbunathan H, et al. Punctuated evolution of canonical genomic aberrations in uveal melanoma. *Nat Commun.* 2018;9:116.
267. Rodrigues M, Mobuchon L, Houy A, et al. Evolutionary Routes in Metastatic Uveal Melanomas Depend on MBD4 Alterations. *Clin Cancer Res.* 2019;25(18):5513-5524.
268. Shain AH, Bagger MM, Yu R, et al. The genetic evolution of metastatic uveal melanoma. *Nat Genet.* 2019;51:1123-1130.
269. Ewens KG, Kanetsky PA, Richards-Yutz J, et al. Chromosome 3 status combined with BAP1 and EIF1AX mutation profiles are associated with metastasis in uveal melanoma. *Invest Ophthalmol Vis Sci.* 2014;55(8):5160-5167.
270. Griewank KG, van de Nes J, Schilling B, et al. Genetic and clinico-pathologic analysis of metastatic uveal melanoma. *Mod Pathol.* 2014;27(2):175-183.

271. Johansson P, Aoude LG, Wadt K, et al. Deep sequencing of uveal melanoma identifies a recurrent mutation in PLCB4. *Oncotarget*. 2016;7(4):4624-4631.
272. Moore AR, Ceraudo E, Sher JJ, et al. Recurrent activating mutations of G-protein-coupled receptor CYSLTR2 in uveal melanoma. *Nat Genet*. 2016;48(6):675-680.
273. Carbone M, Yang H, Pass HI, Krausz T, Testa JR, Gaudino G. BAP1 and cancer. *Nat Rev Cancer*. 2013;13(3):153-159.
274. Furney SJ, Pedersen M, Gentien D, et al. SF3B1 mutations are associated with alternative splicing in uveal melanoma. *Cancer Discov*. 2013;3(10):1122-1129.
275. Turell ME, Tubbs RR, Biscotti CV, Singh AD. Uveal melanoma: prognostication. *Monogr Clin Cytol*. 2012;21:55-60.
276. Onken MD, Worley LA, Tuscan MD, Harbour JW. An accurate, clinically feasible multi-gene expression assay for predicting metastasis in uveal melanoma. *J Mol Diagn*. 2010;12(4):461-468.
277. Field MG, Decatur CL, Kurtenbach S, et al. PRAME as an Independent Biomarker for Metastasis in Uveal Melanoma. *Clin Cancer Res*. 2016;22(5):1234-1242.
278. Augsburger JJ, Correa ZM, Freire J, Brady LW. Long-term survival in choroidal and ciliary body melanoma after enucleation versus plaque radiation therapy. *Ophthalmology*. 1998;105(9):1670-1678.
279. Augsburger JJ, Schneider S, Freire J, Brady LW. Survival following enucleation versus plaque radiotherapy in statistically matched subgroups of patients with choroidal melanomas: results in patients treated between 1980 and 1987. *Graefes Arch Clin Exp Ophthalmol*. 1999;237(7):558-567.
280. Stallard HB. Partial choroidectomy. *Br J Ophthalmol*. 1966;50(11):660-662.
281. Damato BE. Local resection of uveal melanoma. *Dev Ophthalmol*. 2012;49:66-80.
282. Nag S, Quivey JM, Earle JD, Followill D, Fontanesi J, Finger PT. The American Brachytherapy Society recommendations for brachytherapy of uveal melanomas. *Int J Radiat Oncol Biol Phys*. 2003;56(2):544-555.
283. Gragoudas ES, Goitein M, Koehler A, et al. Proton irradiation of choroidal melanomas. Preliminary results. *Arch Ophthalmol*. 1978;96(9):1583-1591.
284. Char DH, Kroll SM, Castro J. Ten-year follow-up of helium ion therapy for uveal melanoma. *Am J Ophthalmol*. 1998;125(1):81-89.
285. Desjardins L, Lumbroso-Le Rouic L, Levy-Gabriel C, et al. Treatment of uveal melanoma by accelerated proton beam. *Dev Ophthalmol*. 2012;49:41-57.
286. Dendale R, Lumbroso-Le Rouic L, Noel G, et al. Proton beam radiotherapy for uveal melanoma: results of Curie Institut-Orsay proton therapy center (ICPO). *Int J Radiat Oncol Biol Phys*. 2006;65(3):780-787.
287. Danish H, Ferris MJ, Balagamwala E, et al. Comparative outcomes and toxicities for ruthenium-106 versus palladium-103 in the treatment of choroidal melanoma. *Melanoma Res*. 2018;28(2):120-125.
288. Gorovets D, Gagne NL, Melhus CS. Dosimetric and radiobiologic comparison of (103)Pd COMS plaque brachytherapy and Gamma Knife radiosurgery for choroidal melanoma. *Brachytherapy*. 2017;16(2):433-443.

289. Abrams MJ, Gagne NL, Melhus CS, Mignano JE. Brachytherapy vs. external beam radiotherapy for choroidal melanoma: Survival and patterns-of-care analyses. *Brachytherapy*. 2016;15:216-223.
290. Seddon JM, Gragoudas ES, Egan KM, et al. Relative survival rates after alternative therapies for uveal melanoma. *Ophthalmology*. 1990;97(6):769-777.
291. Puusaari I, Heikkonen J, Summanen P, Tarkkanen A, Kivela T. Iodine brachytherapy as an alternative to enucleation for large uveal melanomas. *Ophthalmology*. 2003;110(11):2223-2234.
292. Shields CL, Shields JA, Perez N, Singh AD, Cater J. Primary transpupillary thermotherapy for small choroidal melanoma in 256 consecutive cases: outcomes and limitations. *Ophthalmology*. 2002;109(2):225-234.
293. Lommatzsch P. Treatment of choroidal melanomas with 106Ru/106Rh beta-ray applicators. *Surv Ophthalmol*. 1974;19(2):85-100.
294. Lommatzsch PK, Kirsch IH. 106Ru/106Rh plaque radiotherapy for malignant melanomas of the choroid. With follow-up results more than 5 years. *Doc Ophthalmol*. 1988;68(3-4):225-238.
295. Bornfeld N, Chauvel P, Sauerwein W, et al. Metastatic disease, eye retention and visual function in conservative treatment of uveal melanoma. *Front Radiat Ther Oncol*. 1997;30:97-110.
296. Davelaar J, Schaling DF, Hennen LA, Broerse JJ. Dosimetry of ruthenium-106 eye applicators. *Med Phys*. 1992;19(3):691-694.
297. Kleineidam M, Guthoff R, Bentzen SM. Rates of local control, metastasis, and overall survival in patients with posterior uveal melanomas treated with ruthenium-106 plaques. *Radiother Oncol*. 1993;28(2):148-156.
298. Müller RP, Busse H, Pötter R, Kroll P, Haverkamp U. Results of high dose 106-ruthenium irradiation of choroidal melanomas. *Int J Radiat Oncol Biol Phys*. 1986;12(10):1749-1755.
299. Seregard S, aft Trampe E, Lax I, Kock E, Lundell G. Results following episcleral ruthenium plaque radiotherapy for posterior uveal melanoma. The Swedish experience. *Acta Ophthalmol Scand*. 1997;75(1):11-16.
300. Summanen P, Immonen I, Kivelä T, Tommila P, Heikkonen J, Tarkkanen A. Visual outcome of eyes with malignant melanoma of the uvea after ruthenium plaque radiotherapy. *Ophthalmic Surg Lasers*. 1995;26(5):449-460.
301. Summanen P, Immonen I, Heikkonen J, Tommila P, Laatikainen L, Tarkkanen A. Survival of patients and metastatic and local recurrent tumor growth in malignant melanoma of the uvea after ruthenium plaque radiotherapy. *Ophthalmic Surg*. 1993;24(2):82-90.
302. Summanen P, Immonen I, Kivelä T, Tommila P, Heikkonen J, Tarkkanen A. Radiation related complications after ruthenium plaque radiotherapy of uveal melanoma. *Br J Ophthalmol*. 1996;80(8):732-739.
303. Papageorgiou KI, Cohen VM, Bunce C, Kinsella M, Hungerford JL. Predicting local control of choroidal melanomas following (106)Ru plaque brachytherapy. *Br J Ophthalmol*. 2011;95(2):166-170.
304. Stannard C, Sauerwein W, Maree G, Lecuona K. Radiotherapy for ocular tumours. *Eye (Lond)*. 2013;27(2):119-127.

305. Wilkinson DA, Kolar M, Fleming PA, Singh AD. Dosimetric comparison of 106Ru and 125I plaques for treatment of shallow (≤ 5 mm) choroidal melanoma lesions. *Br J Radiol*. 2008;81(970):784-789.
306. Laube T, Flühs D, Kessler C, Bornfeld N. Determination of surgeon's absorbed dose in iodine 125 and ruthenium 106 ophthalmic plaque surgery. *Ophthalmology*. 2000;107(2):366-368; discussion 368-369.
307. Damato B, Patel I, Campbell IR, Mayles HM, Errington RD. Local tumor control after 106Ru brachytherapy of choroidal melanoma. *Int J Radiat Oncol Biol Phys*. 2005;63(2):385-391.
308. Damato B, Patel I, Campbell IR, Mayles HM, Errington RD. Visual acuity after Ruthenium(106) brachytherapy of choroidal melanomas. *Int J Radiat Oncol Biol Phys*. 2005;63(2):392-400.
309. Bergman L, Nilsson B, Lundell G, Lundell M, Seregard S. Ruthenium brachytherapy for uveal melanoma, 1979-2003: survival and functional outcomes in the Swedish population. *Ophthalmology*. 2005;112(5):834-840.
310. Russo A, Laguardia M, Damato B. Eccentric ruthenium plaque radiotherapy of posterior choroidal melanoma. *Graefes Arch Clin Exp Ophthalmol*. 2012;250(10):1533-1540.
311. Wen JC, McCannel TA. Treatment of radiation retinopathy following plaque brachytherapy for choroidal melanoma. *Curr Opin Ophthalmol*. 2009;20(3):200-204.
312. Finger PT. Intraoperative echographic localization of iodine-125 episcleral plaque for brachytherapy of choroidal melanoma. *Am J Ophthalmol*. 2000;130(4):539-540.
313. Geeraerts T, Newcombe VF, Coles JP, et al. Use of T2-weighted magnetic resonance imaging of the optic nerve sheath to detect raised intracranial pressure. *Crit Care*. 2008;12(5):R114.
314. Verschueren KM, Creutzberg CL, Schalijs-Delfos NE, et al. Long-term outcomes of eye-conserving treatment with Ruthenium(106) brachytherapy for choroidal melanoma. *Radiother Oncol*. 2010;95(3):332-338.
315. Wilson MW, Hungerford JL. Comparison of episcleral plaque and proton beam radiation therapy for the treatment of choroidal melanoma. *Ophthalmology*. 1999;106(8):1579-1587.
316. Marinkovic M, Horeweg N, Fiocco M, et al. Ruthenium-106 brachytherapy for choroidal melanoma without transpupillary thermotherapy: Similar efficacy with improved visual outcome. *Eur J Cancer*. 2016;68:106-113.
317. Tarmann L, Wackernagel W, Avian A, et al. Ruthenium-106 plaque brachytherapy for uveal melanoma. *Br J Ophthalmol*. 2015;99(12):1644-1649.
318. Sagoo MS, Shields CL, Mashayekhi A, et al. Plaque radiotherapy for juxtapapillary choroidal melanoma: tumor control in 650 consecutive cases. *Ophthalmology*. 2011;118(2):402-407.
319. Seregard S, Lundell G, Lax I, af Trampe E, Kock E. Tumour cell proliferation after failed ruthenium plaque radiotherapy for posterior uveal melanoma. *Acta Ophthalmol Scand*. 1997;75(2):148-154.
320. Rouberol F, Roy P, Kodjikian L, Gérard JP, Jean-Louis B, Grange JD. Survival, anatomic, and functional long-term results in choroidal and ciliary body melanoma after ruthenium brachytherapy (15 years' experience with beta-rays). *Am J Ophthalmol*. 2004;137(5):893-900.

321. Belaïd A, Nasr C, Jmour O, et al. Brachytherapy of Uveal Melanomas with Ruthenium-106 Plaques. *Asian Pac J Cancer Prev*. 2016;17(12):5281-5285.
322. Rospond-Kubiak I, Wróblewska-Zierhoffer M, Twardosz-Pawlik H, Kocięcki J. Ruthenium brachytherapy for uveal melanoma - single institution experience. *J Contemp Brachytherapy*. 2017;9(6):548-552.
323. Perri P, Fiorica F, D'Angelo S, et al. Ruthenium-106 eye plaque brachytherapy in the conservative treatment of uveal melanoma: a mono-institutional experience. *Eur Rev Med Pharmacol Sci*. 2012;16(14):1919-1924.
324. Archer DB, Amoaku WM, Gardiner TA. Radiation retinopathy--clinical, histopathological, ultrastructural and experimental correlations. *Eye (Lond)*. 1991;5 (Pt 2):239-251.
325. Archer DB, Gardiner TA. Ionizing radiation and the retina. *Curr Opin Ophthalmol*. 1994;5(3):59-65.
326. Krema H, Xu W, Payne D, Vasquez LM, Pavlin CJ, Simpson R. Factors predictive of radiation retinopathy post (125)Iodine brachytherapy for uveal melanoma. *Can J Ophthalmol*. 2011;46(2):158-163.
327. Gupta A, Muecke JS. Treatment of radiation maculopathy with intravitreal injection of bevacizumab (Avastin). *Retina*. 2008;28(7):964-968.
328. Espensen CA, Appelt AL, Fog LS, Gothelf AB, Thariat J, Kiilgaard JF. Predicting Visual Acuity Deterioration and Radiation-Induced Toxicities after Brachytherapy for Choroidal Melanomas. *Cancers*. 2019;11(8).
329. Browne AW, Dandapani SV, Jennelle R, et al. Outcomes of medium choroidal melanomas treated with ruthenium brachytherapy guided by three-dimensional pretreatment modeling. *Brachytherapy*. 2015;14(5):718-725.
330. Finger PT. Tumour location affects the incidence of cataract and retinopathy after ophthalmic plaque radiation therapy. *Br J Ophthalmol*. 2000;84(9):1068-1070.
331. Melia BM, Abramson DH, Albert DM, et al. Collaborative ocular melanoma study (COMS) randomized trial of I-125 brachytherapy for medium choroidal melanoma. I. Visual acuity after 3 years COMS report no. 16. *Ophthalmology*. 2001;108(2):348-366.
332. Chojniak MM, Chojniak R, Nishimoto IN, Allemann N, Erwenne CM. Primary transpupillary thermotherapy for small choroidal melanoma. *Graefes Arch Clin Exp Ophthalmol*. 2011;249(12):1859-1865.
333. Stoffelns BM. Primary transpupillary thermotherapy (TTT) for malignant choroidal melanoma. *Acta Ophthalmol Scand*. 2002;80(1):25-31.
334. Shields CL, Sioufi K, Srinivasan A, et al. Visual Outcome and Millimeter Incremental Risk of Metastasis in 1780 Patients With Small Choroidal Melanoma Managed by Plaque Radiotherapy. *JAMA Ophthalmol*. 2018;136(12):1325-1333.
335. Touth A, Angi M, Dureau S, et al. Long-term visual outcomes for small uveal melanoma staged T1 treated by proton beam radiotherapy. *Cancers*. 2019;11.
336. Pica A, Weber DC, Vallat L, et al. Good long-term visual outcomes of parapapillary choroidal melanoma patients treated with proton therapy: a comparative study. *Int Ophthalmol*. 2020.
337. Journee-de Korver JG, Keunen JE. Thermotherapy in the management of choroidal melanoma. *Prog Retin Eye Res*. 2002;21(3):303-317.

338. Journee-de Korver JG, Oosterhuis JA, de Wolff-Rouendaal D, Kemme H. Histopathological findings in human choroidal melanomas after transpupillary thermotherapy. *Br J Ophthalmol*. 1997;81(3):234-239.
339. Bartlema YM, Oosterhuis JA, Journee-De Korver JG, Tjho-Heslinga RE, Keunen JE. Combined plaque radiotherapy and transpupillary thermotherapy in choroidal melanoma: 5 years' experience. *Br J Ophthalmol*. 2003;87(11):1370-1373.
340. Tjho-Heslinga RE, Davelaar J, Kemme HM, et al. Results of ruthenium irradiation of uveal melanomas: the Dutch experience. *Radiother Oncol*. 1999;53(2):133-137.
341. Journée-De Korver JG, Oosterhuis JA, Kakebeeke-Kemme HM, de Wolff-Rouendaal D. Transpupillary thermotherapy (TTT) by infrared irradiation of choroidal melanoma. *Doc Ophthalmol*. 1992;82(3):185-191.
342. Shields CL, Shields JA, Cater J, et al. Transpupillary thermotherapy for choroidal melanoma: tumor control and visual results in 100 consecutive cases. *Ophthalmology*. 1998;105(4):581-590.
343. Shields CL, Cater J, Shields JA, et al. Combined plaque radiotherapy and transpupillary thermotherapy for choroidal melanoma: tumor control and treatment complications in 270 consecutive patients. *Arch Ophthalmol*. 2002;120(7):933-940.
344. Godfrey DG, Waldron RG, Capone A, Jr. Transpupillary thermotherapy for small choroidal melanoma. *Am J Ophthalmol*. 1999;128(1):88-93.
345. Spire M, Devouassoux MS, Kodjikian L, Janin-Magnificat H, Fleury J, Grange JD. Primary transpupillary thermotherapy for 18 small posterior pole uveal melanomas. *Am J Ophthalmol*. 2006;141(5):840-849.
346. Gragoudas ES, Goitein M, Seddon J, et al. Preliminary results of proton beam irradiation of macular and paramacular melanomas. *Br J Ophthalmol*. 1984;68(7):479-485.
347. Damato B, Kacperek A, Chopra M, Campbell IR, Errington RD. Proton beam radiotherapy of choroidal melanoma: the Liverpool-Clatterbridge experience. *Int J Radiat Oncol Biol Phys*. 2005;62(5):1405-1411.
348. Egger E, Schalenbourg A, Zografos L, et al. Maximizing local tumor control and survival after proton beam radiotherapy of uveal melanoma. *Int J Radiat Oncol Biol Phys*. 2001;51(1):138-147.
349. Gragoudas E, Li W, Goitein M, Lane AM, Munzenrider JE, Egan KM. Evidence-based estimates of outcome in patients irradiated for intraocular melanoma. *Arch Ophthalmol*. 2002;120(12):1665-1671.
350. Seddon JM, Gragoudas ES, Polivogianis L, et al. Visual outcome after proton beam irradiation of uveal melanoma. *Ophthalmology*. 1986;93(5):666-674.
351. Patel AV, Lane AM, Morrison MA, et al. Visual outcomes after proton beam irradiation for choroidal melanomas involving the fovea. *Ophthalmology*. 2016;123:369-377.
352. Gragoudas ES, Li W, Lane AM, Munzenrider J, Egan KM. Risk factors for radiation maculopathy and papillopathy after intraocular irradiation. *Ophthalmology*. 1999;106(8):1571-1577; discussion 1577-1578.
353. Treatment of Age-related macular degeneration with Photodynamic therapy (TAP) study group. Photodynamic therapy of subfoveal choroidal neovascularization in age-related macular degeneration with verteporfin: one-year results of 2 randomized clinical trials -TAP report. *Arch Ophthalmol*. 1999;117:1329-1345.

354. Fabian ID, Stacey AW, Harby LA, Arora AK, Sagoo MS, Cohen VML. Primary photodynamic therapy with verteporfin for pigmented posterior pole cT1a choroidal melanoma: a 3-year retrospective analysis. *Br J Ophthalmol*. 2018;102(12):1705-1710.
355. Turkoglu EB, Pointdujour-Lim R, Mashayekhi A, Shields CL. Photodynamic therapy as primary treatment for small choroidal melanoma. *Retina*. 2019;39(7):1319-1325.
356. Jmor F, Hussain RN, Damato BE, Heimann H. Photodynamic therapy as initial treatment for small choroidal melanomas. *Photodiagnosis Photodyn Ther*. 2017;20:175-181.
357. Fabian ID, Stacey AW, Papastefanou V, et al. Primary photodynamic therapy with verteporfin for small pigmented posterior pole choroidal melanoma. *Eye (Lond)*. 2017;31:519-528.
358. Rundle P. Treatment of posterior uveal melanoma with multi-dose photodynamic therapy. *Br J Ophthalmol*. 2014;98:494-497.
359. Cerman E, Çekiç O. Clinical use of photodynamic therapy in ocular tumors. *Surv Ophthalmol*. 2015;60:557-574.
360. Canal-Fontcuberta I, Salomão DR, Robertson D, et al. Clinical and histopathologic findings after photodynamic therapy of choroidal melanoma. *Retina*. 2012;32:942-948.
361. Grimm J, Scheinberg DA. Will nanotechnology influence targeted cancer therapy? *Semin Radiat Oncol*. 2011;21(2):80-87.
362. Kines RC, Cerio RJ, Roberts JN, et al. Human papillomavirus capsids preferentially bind and infect tumor cells. *Int J Cancer*. 2016;138(4):901-911.
363. Ogawa M, Tomita Y, Nakamura Y, et al. Immunogenic cancer cell death selectively induced by near infrared photoimmunotherapy initiates host tumor immunity. *Oncotarget*. 2017;8(6):10425-10436.
364. Kines RC, Varsavsky I, Choudhary S, et al. An Infrared dye-conjugated virus-like particle for the treatment of primary uveal melanoma. *Mol Cancer Ther*. 2018;17:565-574.
365. Bergman L, Seregard S, Nilsson B, Lundell G, Ringborg U, Ragnarsson-Olding B. Uveal melanoma survival in Sweden from 1960 to 1998. *Invest Ophthalmol Vis Sci*. 2003;44(8):3282-3287.
366. The Collaborative Ocular Melanoma Study Group. Assessment of metastatic disease status at death in 435 patients with large choroidal melanoma in the Collaborative Ocular Melanoma Study (COMS): COMS report no. 15. *Arch Ophthalmol*. 2001;119:670-676.
367. Diener-West M, Reynolds SM, Agugliaro DJ, et al. Screening for metastasis from choroidal melanoma: the Collaborative Ocular Melanoma Study Group Report 23. *J Clin Oncol*. 2004;22(12):2438-2444.
368. Singh AD. Uveal melanoma: implications of tumor doubling time. *Ophthalmology*. 2001;108(5):829-831.
369. Robertson DM. Small choroidal melanomas treated with transpupillary thermotherapy and cryotherapy. *Arch Ophthalmol*. 2008;126(8):1156-1157.
370. Malclès A, Kivelä T, Svetlosakova Z, et al. Small metastasizing choroidal melanomas. *Acta Ophthalmol*. 2015;93(2):e160-166.
371. Thomas JV, Green WR, Maumenee AE. Small choroidal melanomas. A long-term follow-up study. *Arch Ophthalmol*. 1979;97(5):861-864.

372. Sobrin L, Schiffman JC, Markoe AM, Murray TG. Outcomes of iodine 125 plaque radiotherapy after Initial observation of suspected small choroidal melanomas: A pilot study. *Ophthalmology*. 2005;112(10):1777-1783.
373. Barr CC, Sipperley JO, Nicholson DH. Small melanomas of the choroid. *Arch Ophthalmol*. 1978;96(9):1580-1582.
374. Gündüz K, Karslıoğlu MZ, Köse K. Primary transpupillary thermotherapy of choroidal melanocytic lesions. *Middle East Afr J Ophthalmol*. 2011;18(2):183-188.
375. Lane AM, Egan KM, Kim IK, Gragoudas ES. Mortality after diagnosis of small melanocytic lesions of the choroid. *Arch Ophthalmol*. 2010;128(8):996-1000.
376. Eskelin S, Pyrhönen S, Summanen P, Prause JU, Kivelä T. Screening for metastatic malignant melanoma of the uvea revisited. *Cancer*. 1999;85(5):1151-1159.
377. Char DH. Metastatic choroidal melanoma. *Am J Ophthalmol*. 1978;86(1):76-80.
378. Rietschel P, Panageas KS, Hanlon C, Patel A, Abramson DH, Chapman PB. Variates of survival in metastatic uveal melanoma. *J Clin Oncol*. 2005;23(31):8076-8080.
379. Lorigan JG, Wallace S, Mavligit GM. The prevalence and location of metastases from ocular melanoma: imaging study in 110 patients. *AJR Am J Roentgenol*. 1991;157(6):1279-1281.
380. Cerbone L, Van Ginderdeuren R, Van den Oord J, et al. Clinical presentation, pathological features and natural course of metastatic uveal melanoma, an orphan and commonly fatal disease. *Oncology*. 2014;86(3):185-189.
381. Mäkitie T, Kivelä T. Cardiac Metastasis From Uveal Melanoma. *Arch Ophthalmol*. 2001;119(1):139-140.
382. Eskelin S, Kivelä T. Imaging to detect metastases from malignant uveal melanoma. *Arch Ophthalmol*. 2002;120(5):676.
383. Marshall E, Romaniuk C, Ghaneh P, et al. MRI in the detection of hepatic metastases from high-risk uveal melanoma: a prospective study in 188 patients. *Br J Ophthalmol*. 2013;97(2):159-163.
384. Gombos DS, Van Quill KR, Uusitalo M, O'Brien JM. Geographic disparities in diagnostic screening for metastatic uveal melanoma. *Ophthalmology*. 2004;111(12):2254-2258.
385. Francis JH, Patel SP, Gombos DS, Carvajal RD. Surveillance options for patients with uveal melanoma following definitive management. *Am Soc Clin Oncol Educ Book*. 2013:382-387.
386. Rantala ES, Peltola E, Helminen H, Hernberg M, Kivelä TT. Hepatic Ultrasonography Compared with Computed Tomography and Magnetic Resonance Imaging at Diagnosis of Metastatic Uveal Melanoma. *Am J Ophthalmol*. 2020.
387. Khoja L, Atenafu EG, Suci S, et al. Meta-analysis in metastatic uveal melanoma to determine progression free and overall survival benchmarks: an international rare cancers initiative (IRCI) ocular melanoma study. *Ann Oncol*. 2019;30(8):1370-1380.
388. Lane AM, Kim IK, Gragoudas ES. Survival Rates in Patients After Treatment for Metastasis From Uveal Melanoma. *JAMA Ophthalmol*. 2018;136(9):981-986.
389. Augsburger JJ, Corrêa ZM, Shaikh AH. Effectiveness of treatments for metastatic uveal melanoma. *Am J Ophthalmol*. 2009;148(1):119-127.
390. Albert DM, Niffenegger AS, Willson JK. Treatment of metastatic uveal melanoma: review and recommendations. *Surv Ophthalmol*. 1992;36(6):429-438.
391. Aoyama T, Mastrangelo MJ, Berd D, et al. Protracted survival after resection of metastatic uveal melanoma. *Cancer*. 2000;89(7):1561-1568.

392. Fournier GA, Albert DM, Arrigg CA, Cohen AM, Lamping KA, Seddon JM. Resection of solitary metastasis. Approach to palliative treatment of hepatic involvement with choroidal melanoma. *Arch Ophthalmol*. 1984;102(1):80-82.
393. Gündüz K, Shields JA, Shields CL, Sato T, Mastrangelo MJ. Surgical removal of solitary hepatic metastasis from choroidal melanoma. *Am J Ophthalmol*. 1998;125(3):407-409.
394. Frenkel S, Nir I, Hendler K, et al. Long-term survival of uveal melanoma patients after surgery for liver metastases. *Br J Ophthalmol*. 2009;93(8):1042-1046.
395. Hsueh EC, Essner R, Foshag LJ, Ye X, Wang HJ, Morton DL. Prolonged survival after complete resection of metastases from intraocular melanoma. *Cancer*. 2004;100(1):122-129.
396. Pe'er J. Ruthenium-106 brachytherapy. *Dev Ophthalmol*. 2012;49:27-40.
397. Shields CL. The hunt for the secrets of uveal melanoma. *Clin Exp Ophthalmol*. 2008;36(3):277-280.
398. Schwartz M. Biomathematical approach to clinical tumor growth. *Cancer*. 1961;14:1272-1294.
399. Klein C, Bueler H, Mulligan RC. Comparative analysis of genetically modified dendritic cells and tumor cells as therapeutic cancer vaccines. *J Exp Med*. 2000;191:1699-1708.
400. Gass JD. Comparison of uveal melanoma growth rates with mitotic index and mortality. *Arch Ophthalmol*. 1985;103:924-931.
401. Laquis SJ, Freeman JM, Fleming JC, Wilson MW, Haik BG. A rapidly growing choroidal melanoma. *Am J Ophthalmol*. 2002;133:580-581.
402. Harbour JW, Char DH, Kroll S, Quivey JM, Castro J. Metastatic risk for distinct patterns of postirradiation local recurrence of posterior uveal melanoma. *Ophthalmology*. 1997;104:1785-1792; discussion 1792-1783.
403. Puusaari I, Heikkonen J, Kivelä T. Ocular complications after iodine brachytherapy for large uveal melanomas. *Ophthalmology*. 2004;111:1768-1777.
404. Gooley TA, Leisenring W, Crowley J, Storer BE. *Handbook of Statistics in Clinical Oncology*. New York, NY: Marcel Dekker; 2001.
405. Gray RJ. A class of K-sample tests for comparing the cumulative incidence of a competing risk. *Ann Stat*. 1988;16:1141-1154.
406. Kaiserman I, Anteby I, Chowers I, Blumenthal EZ, Kliars I, Pe'er J. Changes in ultrasound findings in posterior uveal melanoma after Ruthenium 106 brachytherapy. *Ophthalmology*. 2002;109(6):1137-1141.
407. Semenova E, Finger PT. Palladium-103 radiation therapy for small choroidal melanoma. *Ophthalmology*. 2013;120(11):2353-2357.
408. Nguyen BT, Kim RS, Bretana ME, Kegley E, Scheffler AC. Association between traditional clinical high-risk features and gene expression profile classification in uveal melanoma. *Graefes Arch Clin Exp Ophthalmol*. 2018;256(2):421-427.